

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
11 April 2002 (11.04.2002)

PCT

(10) International Publication Number  
WO 02/29080 A2

AVAILABLE COPY  
BEST AVAILABLE COPY

(51) International Patent Classification<sup>7</sup>: C12P 13/04,  
13/08, C12N 1/21, 15/11, 15/60 // (C12P 13/04, C12R  
1:19) (C12P 13/08, C12R 1:19)

17, 33602 Bielefeld (DE). HERMANN, Thomas; Zirkon-  
strasse 8, 33739 Bielefeld (DE). THIERBACH, Georg;  
Gunststrasse 21, 33613 Bielefeld (DE).

(21) International Application Number: PCT/EP01/10209

(22) International Filing Date:  
5 September 2001 (05.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
100 48 605.3 30 September 2000 (30.09.2000) DE  
100 55 516.0 9 November 2000 (09.11.2000) DE  
101 30 192.8 22 June 2001 (22.06.2001) DE

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,  
ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

(71) Applicant: DEGUSSA AG [DE/DE]; Bennigsenplatz 1,  
40474 Düsseldorf (DE).

(72) Inventors: RIEPING, Mechthild; Mönkebergstrasse 1,  
33619 Bielefeld (DE). BASTUCK, Christine; Siekerwall

[Continued on next page]

(54) Title: FERMENTATION PROCESS FOR THE PREPARATION OF L-AMINO ACIDS USING STRAINS OF THE FAMILY  
ENTEROBACTERIACEAE



(57) Abstract: The invention relates to a fermentation process for the preparation of L-amino acids, especially L-threonine, in which the following steps are carried out: a) fermentation of the microorganisms of the family Enterobacteriaceae producing the desired L-amino acid, in which microorganisms at least the pckA gene and/or the open reading frames yjfA and ytfP are, individually or jointly, attenuated and, in particular, switched off, b) enrichment of the L-amino acid in the medium or in the bacterial cells, and c) isolation of the L-amino acid.



**Published:**

— without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Fermentation Process for the Preparation of L-Amino Acids  
using Strains of the Family Enterobacteriaceae**

Field of the Invention

The present invention relates to a fermentation process for  
5 the preparation of L-amino acids, especially L-threonine,  
using strains of the family Enterobacteriaceae in which at  
least the pckA gene is attenuated.

Prior Art

L-Amino acids are used in animal nutrition, in human  
10 medicine and in the pharmaceutical industry.

It is known to prepare L-amino acids by the fermentation of  
strains of Enterobacteriaceae, especially *Escherichia coli*  
and *Serratia marcescens*. Because of their great  
importance, attempts are constantly being made to improve  
15 the preparative processes. Improvements to the processes  
may relate to measures involving the fermentation  
technology, e.g. stirring and oxygen supply, or the  
composition of the nutrient media, e.g. the sugar  
concentration during fermentation, or the work-up to the  
20 product form, e.g. by ion exchange chromatography, or the  
intrinsic productivity characteristics of the microorganism  
itself.

The productivity characteristics of these microorganisms  
are improved by using methods of mutagenesis, selection and  
25 mutant choice to give strains which are resistant to  
antimetabolites, e.g. the threonine analogue  $\alpha$ -amino- $\beta$ -  
hydroxyvaleric acid (AHV) or auxotrophic for metabolites of  
regulatory significance, and produce L-amino acids, e.g. L-  
threonine.

30 Methods of recombinant DNA technology have also been used  
for some years to improve L-amino acid-producing strains of

the family Enterobacteriaceae by amplifying individual amino acid biosynthesis genes and studying the effect on production.

#### Object of the Invention

- 5 The object which the inventors set themselves was to provide novel procedures for improving the preparation of L-amino acids, especially L-threonine, by fermentation.

#### Summary of the Invention

- The invention provides a fermentation process for the  
10 preparation of L-amino acids, especially L-threonine, using microorganisms of the family Enterobacteriaceae which, in particular, already produce L-threonine and in which the nucleotide sequence (pckA gene) coding for the enzyme phosphoenolpyruvate carboxykinase (PEP carboxykinase)  
15 (EC 4.1.1.49) is attenuated.

#### Detailed Description of the Invention

- Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-  
20 asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan, L-homoserine and L-arginine. L-Threonine is particularly preferred.

- 25 In this context the term "attenuation" describes the reduction or switching-off, in a microorganism, of the intracellular activity of one or more enzymes (proteins) which are coded for by the appropriate DNA, for example by using a weak promoter or a gene or allele which codes for  
30 an appropriate enzyme with low activity, or inactivating

the appropriate enzyme (protein), and optionally combining these measures.

By attenuation measures, the activity or concentration of the corresponding protein is in general reduced to 0 to 75%, 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild-type protein or of the activity or concentration of the protein in the starting microorganism.

The process is characterized in that the following steps are carried out:

- a) fermentation of microorganisms of the family Enterobacteriaceae in which at least the pckA gene is attenuated,
- b) enrichment of the appropriate L-amino acid in the medium or in the cells of the microorganisms of the family Enterobacteriaceae, and
- c) isolation of the desired L-amino acid.

The microorganisms provided by the present invention can produce L-amino acids from glucose, sucrose, lactose, fructose, maltose, molasses, optionally starch or optionally cellulose, or from glycerol and ethanol. Said microorganisms are representatives of the family Enterobacteriaceae selected from the genera Escherichia, Erwinia, Providencia and Serratia. The genera Escherichia and Serratia are preferred. The species Escherichia coli and Serratia marcescens may be mentioned in particular among the genera Escherichia and Serratia respectively.

Examples of suitable strains, particularly L-threonine-producing strains, of the genus Escherichia, especially of the species Escherichia coli, are:

- Escherichia coli TF427  
Escherichia coli H4578  
Escherichia coli KY10935  
Escherichia coli VNIIGenetika MG442  
5 Escherichia coli VNIIGenetika M1  
Escherichia coli VNIIGenetika 472T23  
Escherichia coli BKIIM B-3996  
Escherichia coli kat 13  
Escherichia coli KCCM-10132.

- 10 Examples of suitable L-threonine-producing strains of the  
genus *Serratia*, especially of the species *Serratia*  
*marcescens*, are:

- Serratia marcescens HNr21  
Serratia marcescens TLR156  
15 Serratia marcescens T2000.

- L-Threonine-producing strains of the family  
Enterobacteriaceae preferably possess, inter alia, one or  
more genetic or phenotypic characteristics selected from  
the group comprising resistance to  $\alpha$ -amino- $\beta$ -hydroxyvaleric  
20 acid, resistance to thialysine, resistance to ethionine,  
resistance to  $\alpha$ -methylserine, resistance to diaminosuccinic  
acid, resistance to  $\alpha$ -aminobutyric acid, resistance to  
borrelidine, resistance to rifampicin, resistance to valine  
analogues such as valine hydroxamate, resistance to purine  
25 analogues such as 6-dimethylaminopurine, need for L-  
methionine, optionally partial and compensable need for L-  
isoleucine, need for meso-diaminopimelic acid, auxotrophy  
in respect of threonine-containing dipeptides, resistance  
to L-threonine, resistance to L-homoserine, resistance to  
30 L-lysine, resistance to L-methionine, resistance to L-  
glutamic acid, resistance to L-aspartate, resistance to L-  
leucine, resistance to L-phenylalanine, resistance to L-  
serine, resistance to L-cysteine, resistance to L-valine,  
sensitivity to fluoropyruvate, defective threonine

dehydrogenase, optionally capability for sucrose utilization, amplification of the threonine operon, amplification of homoserine dehydrogenase I-aspartate kinase I, preferably of the feedback-resistant form, 5 amplification of homoserine kinase, amplification of threonine synthase, amplification of aspartate kinase, optionally of the feedback-resistant form, amplification of aspartate semialdehyde dehydrogenase, amplification of phosphoenolpyruvate carboxylase, optionally of the 10 feedback-resistant form, amplification of phosphoenolpyruvate synthase, amplification of transhydrogenase, amplification of the RhtB gene product, amplification of the RhtC gene product, amplification of the YfiK gene product, amplification of malate quinone 15 oxidoreductase and amplification of a pyruvate carboxylase and attenuation of acetic acid formation.

It has been found that the production of L-amino acids, especially L-threonine, by microorganisms of the family Enterobacteriaceae is improved after attenuation and, in 20 particular, switching-off of the pckA gene coding for PEP carboxykinase (EC 4.1.1.49).

The nucleotide sequence of the pckA gene of Escherichia coli has been published by Medina et al. (Journal of Bacteriology 172, 7151-7156 (1990)) and can also be taken 25 from the genome sequence of Escherichia coli published by Blattner et al. (Science 277, 1453-1462 (1997)). The nucleotide sequence of the pckA gene of Escherichia coli is represented in SEQ ID No. 1 and the amino acid sequence of the corresponding gene product is represented in SEQ ID No. 30 2.

The pckA genes described in the above literature references can be used according to the invention. It is also possible to use alleles of the pckA gene which result from

the degeneracy of the genetic code or from neutral sense mutations.

Attenuation can be achieved for example by reducing or switching off the expression of the *pckA* gene or the catalytic properties of the enzyme protein. Both measures may optionally be combined.

Gene expression can be reduced by an appropriate culture technique, by genetic modification (mutation) of the signal structures of gene expression, or by means of antisense RNA technology. Examples of signal structures of gene expression are repressor genes, activator genes, operators, promoters, attenuators, ribosome binding sites, the start codon and terminators. Those skilled in the art will find relevant information inter alia in e.g. Jensen and Hammer (Biotechnology and Bioengineering 58, 191-195 (1998)), Carrier and Keasling (Biotechnology Progress 15, 58-64 (1999)), Franch and Gerdes (Current Opinion in Microbiology 3, 159-164 (2000)) and well-known textbooks on genetics and molecular biology, for example the textbook by Knippers ("Molekulare Genetik" ("Molecular Genetics"), 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995) or the textbook by Winnacker ("Gene und Klone" ("From Genes to Clones"), VCH Verlagsgesellschaft, Weinheim, Germany, 1990).

Mutations which cause a change or reduction in the catalytic properties of enzyme proteins are known from the state of the art. Examples which may be mentioned are the studies of Qiu and Goodman (Journal of Biological Chemistry 272, 8611-8617 (1997)), Yano et al. (Proceedings of the National Academy of Sciences USA 95, 5511-5515 (1998)) and Wenthe and Schachmann (Journal of Biological Chemistry 266, 20833-20839 (1991)). Surveys can be found in well-known textbooks on genetics and molecular biology, e.g. the



textbook by Hagemann ("Allgemeine Genetik" ("General Genetics")), Gustav Fischer Verlag, Stuttgart, 1986).

Mutations to be taken into consideration are transitions, transversions, insertions and deletions. Depending on the effect of amino acid exchange on the enzyme activity, the term missense mutations or nonsense mutations is used. Insertions or deletions of at least one base pair in a gene cause frame shift mutations, the result of which is that false amino acids are incorporated or translation is terminated prematurely. Deletions of several codons typically lead to a complete loss of enzyme activity. Instructions for the production of such mutations form part of the state of the art and can be found in well-known textbooks on genetics and molecular biology, e.g. the textbook by Knippers ("Molekulare Genetik" ("Molecular Genetics")), 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995), the textbook by Winnacker ("Gene und Klone" ("From Genes to Clones")), VCH Verlagsgesellschaft, Weinheim, Germany, 1990) or the textbook by Hagemann ("Allgemeine Genetik" ("General Genetics")), Gustav Fischer Verlag, Stuttgart, 1986).

An example of a plasmid by means of which the pckA gene of Escherichia coli can be attenuated and, in particular, switched off by position-specific mutagenesis is plasmid pMAK705 $\Delta$ pckA (Figure 1). It contains only part of the 5' region and part of the 3' region of the pckA gene. A 349 bp segment of the coding region is missing (deletion). The sequence of this DNA, which can be used for mutagenesis of the pckA gene, is represented in SEQ ID No. 3.

The deletion mutation of the pckA gene can be incorporated into suitable strains by gene or allele exchange.

A common method is the method of gene exchange using a conditionally replicating pSC101 derivative, pMAK705, as

described by Hamilton et al. (Journal of Bacteriology 174, 4617-4622 (1989)). Other methods described in the state of the art, for example that of Martinez-Morales et al. (Journal of Bacteriology, 7143-7148 (1999)) or that of Boyd  
5 et al. (Journal of Bacteriology 182, 842-847 (2000)), can also be used.

When exchange has been carried out, the form of the  $\Delta pckA$  allele represented in SEQ ID No. 4, which is a further subject of the invention, is present in the strain in  
10 question.

Mutations in the *pckA* gene or mutations involving expression of the *pckA* gene can also be transferred to different strains by conjugation or transduction.

Furthermore, for the production of L-amino acids,  
15 especially L-threonine, with strains of the family Enterobacteriaceae, it can be advantageous not only to attenuate the *pckA* gene but also to amplify one or more enzymes of the known threonine biosynthetic pathway, or enzymes of the anaplerotic metabolism, or enzymes for the  
20 production of reduced nicotinamide adenine dinucleotide phosphate.

In this context the term "amplification" describes the increase in the intracellular activity, in a microorganism, of one or more enzymes or proteins which are coded for by  
25 the appropriate DNA, for example by increasing the copy number of the gene(s), using a strong promoter or using a gene coding for an appropriate enzyme or protein with a high activity, and optionally combining these measures.

By amplification measures, in particular over-expression,  
30 the activity or concentration of the corresponding protein is in general increased by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on that of the wild-type protein or

the activity or concentration of the protein in the starting microorganism.

Thus, for example, one or more genes selected from the group comprising:

- 5     • the thrABC operon coding for aspartate kinase, homoserine dehydrogenase, homoserine kinase and threonine synthase (US-A-4,278,765),
- the pyc gene coding for pyruvate carboxylase (DE-A-19 831 609),
- 10    • the pps<sub>1</sub> gene coding for phosphoenolpyruvate synthase (Molecular and General Genetics 231, 332 (1992)),
- the ppc gene coding for phosphoenolpyruvate carboxylase (Gene 31, 279-283 (1984)),
- the pntA and pntB genes coding for transhydrogenase  
15       (European Journal of Biochemistry 158, 647-653 (1986)),
- the rhtB gene for homoserine resistance (EP-A-0994190), and
- the rhtC gene for threonine resistance (EP-A-1013765),
- the gdhA gene coding for glutamate dehydrogenase (Gene  
20       27:193-199 (1984))

can be simultaneously amplified and, in particular, overexpressed.

Furthermore, for the production of L-amino acids, especially L-threonine, it can be advantageous not only to  
25   attenuate the pckA gene but also to attenuate and, in particular, switch off one or more genes selected from the group comprising:

- the tdh gene coding for threonine dehydrogenase  
(Ravnikar and Somerville, Journal of Bacteriology 169,  
4716-4721 (1987)),
- 5     • the mdh gene coding for malate dehydrogenase  
      (EC 1.1.1.37) (Vogel et al., Archives in Microbiology  
      149, 36-42 (1987)),
- 10    • the gene product of the open reading frame (orf) yjfa  
      (Accession Number AAC77180 of the National Center for  
      Biotechnology Information (NCBI, Bethesda, MD, USA) and  
      SEQ ID No. 5), and
- the gene-product of the open reading frame (orf) ytfP  
      (Accession Number AAC77179 of the National Center for  
      Biotechnology Information (NCBI, Bethesda, MD, USA) and  
      SEQ ID No. 5),
- 15    or to reduce the expression.

It is preferred to attenuate the open reading frame yjfa  
and/or the open reading frame ytfP.

- It is also possible according to the invention to attenuate  
the open reading frames yjfa and/or ytfP independently of  
20   the pckA gene, in order to achieve an improvement in the  
     amino acids, in particular L-threonine production.

The invention accordingly also provides a process,  
characterized in that the following steps are carried out:

- 25    d)   fermentation of microorganisms of the  
      Enterobacteriaceae family in which at least the  
      open reading frame yjfa and/or ytfP is attenuated,
- e)   enrichment of the L-amino acid in the medium or in  
          the cells of the microorganisms of the  
          Enterobacteriaceae family, and

- f) isolation of the L-threonine, constituents of the fermentation broth and the biomass in its entirety or portions thereof optionally being isolated as a solid product together with the L-amino acid.
- 5 An example of a plasmid by means of which the open reading frames yjfa and ytfP of Escherichia coli can be attenuated and, in particular, switched off by position-specific mutagenesis is plasmid pMAK705 $\Delta$ yjfa (Figure 2). It contains only the 5' and 3' flanks of the ytfP-yjfa region, including very short residues of the open reading frames yjfa and ytfP. A 337 bp long part of the ytfP-yjfa region is missing (deletion). The sequence of this DNA, which can be used for mutagenesis of the ytfP-yjfa region, is represented in SEQ ID No. 6.
- 10
- 15 An further example of a plasmid by means of which the open reading frames yjfa and ytfP of Escherichia coli can be attenuated and, in particular, switched off by position-specific mutagenesis is the plasmid pMAK705 $\Delta$ 90bp (Figure 5). It also contains only the 5' and 3' flanks of the ytfP-yjfa region including very short residues of the open reading frames yjfa and ytfP. A 90 bp long part of the ytfP-yjfa region is missing (deletion). The sequence of this DNA, which can be used for mutagenesis of the ytfP-yjfa region, is represented in SEQ ID No. 7.
- 20
- 25 This deletion mutation can be incorporated into suitable strains by gene or allele replacement. It is also possible to transfer mutations in the open reading frames yjfa and/or ytfP or mutations affecting expression of these open reading frames into various strains by conjugation or transduction.
- 30

When replacement has been carried out, the form of the  $\Delta$ ytfP and  $\Delta$ yjfa allele represented in SEQ ID No. 6 or SEQ

ID No. 7, which are a further subject of the invention, is present in the strain in question.

Furthermore, for the production of L-amino acids, especially L-threonine, it can be advantageous, in addition  
5 to the individual or joint attenuation of the *pckA* gene or of the open reading frames *yjfa* and/or *ytfP*, to switch off undesired secondary reactions (Nakayama: "Breeding of Amino Acid Producing Microorganisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.),  
10 Academic Press, London, UK, 1982).

The microorganisms prepared according to the invention can be cultivated by the batch process or the fed batch process. A summary of known cultivation methods is provided in the textbook by Chmiel (Bioprozesstechnik 1.  
15 Einführung in die Bioverfahrenstechnik (Bioprocess Technology 1. Introduction to Bioengineering) (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen (Bioreactors and Peripheral Equipment) (Vieweg Verlag,  
20 Brunswick/Wiesbaden, 1994)).

The culture medium to be used must appropriately meet the demands of the particular strains. Descriptions of culture media for various microorganisms can be found in the  
25 handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington DC, USA, 1981).

Carbon sources which can be used are sugars and carbohydrates, e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and optionally cellulose, oils  
30 and fats, e.g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, e.g. palmitic acid, stearic acid and linoleic acid, alcohols, e.g. glycerol and ethanol, and

organic acids, e.g. acetic acid. These substances can be used individually or as a mixture.

Nitrogen sources which can be used are organic nitrogen-containing compounds such as peptones, yeast extract, meat  
5 extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate. The nitrogen sources can be used individually or as a mixture.

10 Phosphorus sources which can be used are phosphoric acid, potassium dihydrogenphosphate or dipotassium hydrogenphosphate or the corresponding sodium salts. The culture medium must also contain metal salts, e.g. magnesium sulfate or iron sulfate, which are necessary for  
15 growth. Finally, essential growth-promoting substances such as amino acids and vitamins can be used in addition to the substances mentioned above. Suitable precursors can also be added to the culture medium. Said feed materials can be added to the culture all at once or fed in  
20 appropriately during cultivation.

The pH of the culture is controlled by the appropriate use of basic compounds such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds  
25 such as phosphoric acid or sulfuric acid. Foaming can be controlled using antifoams such as fatty acid polyglycol esters. The stability of plasmids can be maintained by adding suitable selectively acting substances, e.g. antibiotics, to the medium. Aerobic conditions are maintained by introducing oxygen or oxygen-containing  
30 gaseous mixtures, e.g. air, into the culture. The temperature of the culture is normally 25°C to 45°C and preferably 30°C to 40°C. The culture is continued until the formation of L-amino acids or L-threonine has reached a

maximum. This objective is normally achieved within 10 hours to 160 hours.

L-Amino acids can be analyzed by means of anion exchange chromatography followed by ninhydrin derivation, as

- 5 described by Spackman et al. (Analytical Chemistry 30, 1190 (1958)), or by reversed phase HPLC, as described by Lindroth et al. (Analytical Chemistry 51, 1167-1174 (1979)).

- 10 A pure culture of the Escherichia coli K-12 strain DH5 $\alpha$ /pMAK705 was deposited on 12th September 2000 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty as DSM 13720.

- 15 A pure culture of the Escherichia coli K-12 strain MG442 $\Delta$ pckA was deposited on 2nd October 2000 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the  
20 Budapest Treaty as DSM 13761.

- A pure culture of the Escherichia coli K-12 strain B-3996kur $\Delta$ tdh $\Delta$ pckA/pVIC40 was deposited on 9th March 2001 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell  
25 Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty as DSM 14150.

- A pure culture of the Escherichia coli K-12 strain MG442 $\Delta$ 90yjfA was deposited on 9th May 2001 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ =  
30 German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty as DSM 14289.



It is also possible according to the invention individually to attenuate the open reading frames ytfP and yjfA in order to improve the production of L-amino acids.

5 The process according to the invention is used for the preparation of L-amino acids, e.g. L-threonine, L-isoleucine, L-methionine, L-homoserine and L-lysine, especially L-threonine, by fermentation.

The present invention is illustrated in greater detail below with the aid of Examples.

10 The isolation of plasmid DNA from Escherichia coli and all the techniques for restriction, Klenow treatment and alkaline phosphatase treatment were carried out as described by Sambrook et al. (Molecular cloning - A laboratory manual (1989), Cold Spring Harbor Laboratory Press).  
15 Unless indicated otherwise, the transformation of Escherichia coli was carried out as described by Chung et al. (Proceedings of the National Academy of Sciences USA 86, 2172-2175 (1989)).

20 The incubation temperature for the preparation of strains and transformants was 37°C. Temperatures of 30°C and 44°C were used in the gene exchange process of Hamilton et al.

#### Example 1

Construction of the deletion mutation of the pckA gene

25 Parts of the 5' and 3' regions of the pckA gene of Escherichia coli K12 were amplified using the polymerase chain reaction (PCR) and synthetic oligonucleotides. The nucleotide sequence of the pckA gene in E. coli K12 MG1655 (SEQ ID No. 1) was used to synthesize the following PCR primers (MWG Biotech, Ebersberg, Germany):

30 pckA'5'-1: 5' - GATCCGAGCCTGACAGGTTA - 3'  
pckA'5'-2: 5' - GCATGCGCTCGGTCAGGTTA - 3'

pckA'3'-1: 5' - AGGCCTGAAGATGGCACTATCG - 3'

pckA'3'-2: 5' - CCGGAGAAGCGTAGGTGTTA - 3'.

The chromosomal *E. coli* K12 MG1655 DNA used for the PCR was isolated with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany) according to the manufacturer's instructions. An approx. 500 bp DNA fragment from the 5' region of the pckA gene (denoted as pck1) and an approx. 600 bp DNA fragment from the 3' region of the pckA gene (denoted as pck2) could be amplified with the specific primers under standard PCR conditions (Innis et al. (1990), PCR Protocols. A Guide to Methods and Applications, Academic Press) using Taq DNA polymerase (Gibco-BRL, Eggenstein, Germany). The PCR products were each ligated with vector pCR2.1TOPO (TOPO TA Cloning Kit, Invitrogen, Groningen, The Netherlands) according to the manufacturer's instructions and transformed into *E. coli* strain TOP10F'. Plasmid-carrying cells were selected on LB agar containing 50 µg/ml of ampicillin. After isolation of the plasmid DNA, vector pCR2.1TOPOpck2 was cleaved with the restriction enzymes StuI and XbaI and, after separation in 0.8% agarose gel, the pck2 fragment was isolated with the aid of the QIAquick Gel Extraction Kit (QIAGEN, Hilden, Germany). After isolation of the plasmid DNA, vector pCR2.1TOPOpck1 was cleaved with the enzymes EcoRV and XbaI and ligated to the isolated pck2 fragment. The *E. coli* strain DH5α was transformed with the ligation mixture and plasmid-carrying cells were selected on LB agar containing 50 µg/ml of ampicillin. After isolation of the plasmid DNA, control cleavage with the enzymes SpeI and XbaI was used to detect plasmids containing, in cloned form, the mutagenic DNA sequence represented in SEQ ID No. 3. One of the plasmids was denoted as pCR2.1TOPOΔpckA.

Example 2Construction of exchange vector pMAK705 $\Delta$ pckA

After restriction with the enzymes SpeI and XbaI and separation in 0.8% agarose gel, the pckA allele described in Example 1 was isolated from vector pCR2.1TOPO $\Delta$ pckA and ligated to plasmid pMAK705 (Hamilton et al., Journal of Bacteriology 174, 4617-4622 (1989)) which had been digested with the enzyme XbaI. DH5 $\alpha$  was transformed with the ligation mixture and plasmid-carrying cells were selected on LB agar containing 20  $\mu$ g/ml of chloramphenicol. After isolation of the plasmid DNA and cleavage with the enzymes HpaI, KpnI, HindIII, SalI and PstI, successful cloning was detected. The exchange vector formed, pMAK705 $\Delta$ pckA (= pMAK705 $\Delta$ tapckA), is shown in Figure 1.

15 Example 3

Position-specific mutagenesis of the pckA gene in the E. coli strain MG442

The L-threonine-producing E. coli strain MG442 is described in patent US-A-4,278,765 and deposited in the Russian National Collection of Industrial Microorganisms (VKPM, Moscow, Russia) as CMIM B-1628.

The strain MG442 has a resistance to  $\alpha$ -amino- $\beta$ -hydroxyvaleric acid and has an optionally partial and compensable need for L-isoleucine.

25 For exchange of the chromosomal pckA gene for the plasmid-coded deletion construct, MG442 was transformed with plasmid pMAK705 $\Delta$ pckA. The gene exchange was carried out by the selection method described by Hamilton et al. (Journal of Bacteriology 174, 4617-4622 (1989)) and was verified by  
30 standard PCR methods (Innis et al. (1990), PCR Protocols. A

Guide to Methods and Applications, Academic Press) using the following oligonucleotide primers:

pckA'5'-1: 5' - GATCCGAGCCTGACAGGTTA - 3'

pckA'3'-2: 5' - CCGGAGAAGCGTAGGTGTTA - 3'

- 5 The strain obtained was denoted as MG442ΔpckA.

#### Example 4

Preparation of L-threonine with the strain MG442ΔpckA

- MG442ΔpckA was cultivated on minimum medium of the following composition: 3.5 g/l of  $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$ , 1.5 g/l of  $\text{KH}_2\text{PO}_4$ , 1 g/l of  $\text{NH}_4\text{Cl}$ , 0.1 g/l of  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 2 g/l of glucose and 20 g/l of agar. The formation of L-threonine was checked in 10 ml batch cultures contained in 100 ml Erlenmeyer flasks. These were inoculated with 10 ml of a preculture medium of the following composition: 2 g/l of yeast extract, 10 g/l of  $(\text{NH}_4)_2\text{SO}_4$ , 1 g/l of  $\text{KH}_2\text{PO}_4$ , 0.5 g/l of  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 15 g/l of  $\text{CaCO}_3$  and 20 g/l of glucose, and incubated for 16 hours at 37°C and 180 rpm on an ESR incubator from Kühner AG (Birsfelden, Switzerland). 250 μl of this preculture were transferred to 10 ml of a production medium (25 g/l of  $(\text{NH}_4)_2\text{SO}_4$ , 2 g/l of  $\text{KH}_2\text{PO}_4$ , 1 g/l of  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.03 g/l of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.018 g/l of  $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ , 30 g/l of  $\text{CaCO}_3$ , 20 g/l of glucose) and incubated for 48 hours at 37°C. After incubation, the optical density (OD) of the culture suspension was determined with an LP2W photometer from Dr. Lange (Berlin, Germany) at a measurement wavelength of 660 nm.

- The concentration of L-threonine formed was then determined in the sterile-filtered culture supernatant with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by means of ion exchange chromatography and postcolumn reaction with ninhydrin detection.

The result of the experiment is shown in Table 1.

Table 1

Strain	OD (660 nm)	L-Threonine g/l
MG442	6.0	1.5
MG442 $\Delta$ pckA	5.4	3.7

Example 5

- 5 Preparation of L-threonine with the strain  
MG442 $\Delta$ pckA/pMW218gdhA

5.1 Amplification and cloning of the *gdhA* gene

The glutamate dehydrogenase gene from *Escherichia coli* K12 is amplified using the polymerase chain reaction (PCR) and  
10 synthetic oligonucleotides. Starting from the nucleotide  
sequence for the *gdhA* gene in *E. coli* K12 MG1655 (gene  
library: Accession No. AE000270 and No. AE000271) PCR  
primers are synthesized (MWG Biotech, Ebersberg, Germany):

- Gdh1: 5' - TGAACACTTCTGGCGGTACG - 3'  
15 Gdh2: 5' - CCTCGGCGAAGCTAATATGG - 3'

The chromosomal *E. coli* K12 MG1655 DNA employed for the PCR is isolated according to the manufacturers instructions with "QIAGEN Genomic-tips 100/G" (QIAGEN, Hilden, Germany). A DNA fragment approx. 2150 bp in size, which comprises the  
20 *gdhA* coding region and approx. 350 bp 5'-flanking and  
approx. 450 bp 3'-flanking sequences, can be amplified with  
the specific primers under standard PCR conditions (Innis  
et al.: PCR Protocols. A Guide to Methods and Applications,  
1990, Academic Press) with the Pfu-DNA polymerase (Promega  
25 Corporation, Madison, USA). The PCR product is cloned in

the plasmid pCR2.1TOPO and transformed in the E. coli strain TOP10 (Invitrogen, Leek, The Netherlands, Product Description TOPO TA Cloning Kit, Cat. No. K4500-01). Successful cloning is demonstrated by cleavage of the

5 plasmid pCR2.1TOPOgdhA with the restriction enzymes EcoRI and EcoRV. For this, the plasmid DNA is isolated by means of the "QIAprep Spin Plasmid Kits" (QIAGEN, Hilden, Germany) and, after cleavage, separated in a 0.8 % agarose gel.

10 5.2 Cloning of the gdhA gene in the plasmid vector pMW218

The plasmid pCR2.1TOPOgdhA is cleaved with the enzyme EcoRI, the cleavage batch is separated on 0.8% agarose gel and the gdhA fragment 2.1 kbp in size is isolated with the aid of the "QIAquick Gel Extraction Kit" (QIAGEN, Hilden,

15 Germany). The plasmid pMW218 (Nippon Gene, Toyama, Japan) is cleaved with the enzyme EcoRI and ligated with the gdhA fragment. The E. coli strain DH5 $\alpha$  is transformed with the ligation batch and pMW218-carrying cells are selected by plating out on LB agar (Lennox, Virology 1955, 1: 190), to

20 which 20  $\mu$ g/ml kanamycin are added.

Successful cloning of the gdhA gene can be demonstrated after plasmid DNA isolation and control cleavage with EcoRI and EcoRV. The plasmid is called pMW218gdhA (Figure 3).

5.3 Preparation of the strain MG442 $\Delta$ pckA/pMW218gdhA

25 The strain MG442 $\Delta$ pckA obtained in Example 3 and the strain MG442 are transformed with the plasmid pMW218gdhA and transformants are selected on LB agar, which is supplemented with 20  $\mu$ g/ml kanamycin. The strains MG442 $\Delta$ pckA/pMW218gdhA and MG442/pMW218gdhA are formed in

30 this manner.

#### 5.4 Preparation of L-threonine

The preparation of L-threonine by the strains MG442 $\Delta$ pckA/pMW218gdhA and MG442/pMW218gdhA is tested as described in Example 4. The minimal medium and the  
5 preculture medium are additionally supplemented with 20  $\mu$ g/ml kanamycin.

The result of the experiment is summarized in Table 2.

Table 2

Strain	OD (660 nm)	L-Threonine g/l
MG442	6.0	1.5
MG442 $\Delta$ pckA	5.4	3.7
MG442/pMW218gdhA	5.6	2.6
MG442 $\Delta$ pckA/pMW218gdhA	5.5	4.0

#### 10 Example 6

Preparation of L-threonine with the strain MG442 $\Delta$ pckA/pMW219rhtC

##### 6.1 Amplification of the rhtC gene

The rhtC gene from Escherichia coli K12 is amplified using  
15 the polymerase chain reaction (PCR) and synthetic oligonucleotides. Starting from the nucleotide sequence for the rhtC gene in E. coli K12 MG1655 (gene library: Accession No. AE000458, Zakataeva et al. (FEBS Letters 452, 228-232 (1999)), PCR primers are synthesized (MWG Biotech,  
20 Ebersberg, Germany):

RhtC1: 5' - CTGTTAGCATCGGCGAGGCA - 3'

RhtC2: 5' - GCATGTTGATGGCGATGACG - 3'

The chromosomal *E. coli* K12 MG1655 DNA employed for the PCR is isolated according to the manufacturers instructions  
5 with "QIAGEN Genomic-tips 100/G" (QIAGEN, Hilden, Germany). A DNA fragment approx. 800 bp in size can be amplified with the specific primers under standard PCR conditions (Innis et al.: PCR Protocols. A Guide to Methods and Applications, 1990, Academic Press) with Pfu-DNA polymerase (Promega  
10 Corporation, Madison, USA).

#### 6.2 Cloning of the *rhtC* gene in the plasmid vector pMW219

The plasmid pMW219 (Nippon Gene, Toyama, Japan) is cleaved with the enzyme *SamI* and ligated with the *rhtC*-PCR  
15 fragment. The *E. coli* strain DH5 $\alpha$  is transformed with the ligation batch and pMW219-carrying cells are selected on LB agar, which is supplemented with 20  $\mu$ g/ml kanamycin. Successful cloning can be demonstrated after plasmid DNA isolation and control cleavage with *KpnI*, *HindIII* and *NcoI*. The plasmid pMW219*rhtC* is shown in Figure 4.

#### 20 6.3 Preparation of the strain MG442 $\Delta$ pckA/pMW219*rhtC*

The strain MG442 $\Delta$ pckA obtained in Example 3 and the strain MG442 are transformed with the plasmid pMW219*rhtC* and transformants are selected on LB agar, which is  
25 supplemented with 20  $\mu$ g/ml kanamycin. The strains MG442 $\Delta$ pckA/pMW219*rhtC* and MG442/pMW219*rhtC* are formed in this manner.

#### 6.4 Preparation of L-threonine

The preparation of L-threonine by the strains MG442 $\Delta$ pckA/pMW219*rhtC* and MG442/pMW219*rhtC* is tested as  
30 described in Example 4. The minimal medium and the



preculture medium are additionally supplemented with 20 µg/ml kanamycin.

The result of the experiment is summarized in Table 3.

Table 3

Strain	OD (660 nm)	L-Threonine g/l
MG442	6.0	1.5
MG442ΔpckA	5.4	3.7
MG442/pMW219rhtC	5.2	2.9
MG442ΔpckA/pMW219rhtC	4.8	4.4

5

#### Example 7

Preparation of L-threonine with the strain  
B-3996kurΔtdhΔpckA/pVIC40

10 The L-threonine-producing E. coli strain B-3996 is described in US-A- 5,175,107 and deposited at the Russian National Collection for Industrial Microorganisms (VKPM, Moscow, Russia).

15 The strain B-3996 has, inter alia, a resistance to α-amino-β-hydroxyvaleric acid, has an attenuated, in particular switched-off, or defective threonine dehydrogenase, has an enhanced homoserine dehydrogenase I aspartate kinase I in the feed back resistant form, has an optionally partial and compensable need for L-isoleucine and has the ability to utilize sucrose.

### 7.1 Preparation of the strain B-3996kur $\Delta$ tdh $\Delta$ pckA/pVIC40

After culture in antibiotic-free complete medium for approximately ten generations, a derivative of strain B-3996 which no longer contains the plasmid pVIC40 is  
5 isolated. The strain formed is streptomycin-sensitive and is designated B-3996kur.

The method described by Hamilton et al. (Journal of Bacteriology (1989) 171: 4617-4622), which is based on the use of the plasmid pMAK705 with a temperature-sensitive  
10 replicon, was used for incorporation of a deletion into the tdh gene. The plasmid pDR121 (Ravnikar and Somerville, Journal of Bacteriology (1987) 169:4716-4721) contains a DNA fragment from E. coli 3.7 kilo-base pairs (kbp) in size, on which the tdh gene is coded. To generate a  
15 deletion of the tdh gene region, pDR121 is cleaved with the restriction enzymes ClaI and EcoRV and the DNA fragment 5 kbp in size isolated is ligated, after treatment with Klenow enzyme. The ligation batch is transformed in the E. coli strain DH5 $\alpha$  and plasmid-carrying cells are selected on  
20 LB agar, to which 50  $\mu$ g/ml ampicillin are added.

Successful deletion of the tdh gene can be demonstrated after plasmid DNA isolation and control cleavage with EcoRI. The EcoRI fragment 1.7 kbp in size is isolated, and ligated with the plasmid pMAK705, which is partly digested  
25 with EcoRI. The ligation batch is transformed in DH5 $\alpha$  and plasmid-carrying cells are selected on LB agar, to which 20  $\mu$ g/ml chloramphenicol are added. Successful cloning is demonstrated after isolation of the plasmid DNA and cleavage with EcoRI. The pMAK705 derivative formed is  
30 designated pDM32.

For the gene replacement, B-3996kur is transformed with the plasmid pDM32. The replacement of the chromosomal tdh gene with the plasmid-coded deletion construct is carried out by

the selection process described by Hamilton et al. and is verified by standard PCR methods (Innis et al. (1990), PCR Protocols. A Guide to Methods and Applications, Academic Press) with the following oligonucleotide primers:

- 5 Tdh1: 5'-TCGCGACCTATAAGTTTGGG-3'  
Tdh2: 5'-AATACCAGCCCTTGTTTCGTG-3'.

The strain formed is tested for kanamycin sensitivity and is designated B-3996kur $\Delta$ tdh.

- 10 For the position-specific mutagenesis of the pckA gene, B-3996kur $\Delta$ tdh is transformed with the replacement vector pMAK705 $\Delta$ pckA described in Example 2. The replacement of the chromosomal pckA gene by the plasmid-coded deletion construct is carried out as described in Example 3. The strain obtained is called B-3996kur $\Delta$ tdh $\Delta$ pckA.

- 15 B-3996kur $\Delta$ tdh and B-3996kur $\Delta$ tdh $\Delta$ pckA are transformed with the plasmid pVIC40 isolated from B-3996 and plasmid-carrying cells are selected on LB agar with 20  $\mu$ g/ml streptomycin. In each case a selected individual colony is called B-3996kur $\Delta$ tdh/pVIC40 and B-3996kur $\Delta$ tdh $\Delta$ pckA/pVIC40.

- 20 7.2 Preparation of L-threonine

- The preparation of L-threonine by the strains B-3996kur $\Delta$ tdh/pVIC40 and B-3996kur $\Delta$ tdh $\Delta$ pckA/pVIC40 is tested as described in Example 4. The minimal medium, the preculture medium and the production medium are  
25 additionally supplemented with 20  $\mu$ g/ml streptomycin.

The result of the experiment is summarized in Table 4.

Table 4

Strain	OD (660 nm)	L-Threonine g/l
B-3996kur $\Delta$ tdh/pVIC40	4.7	6.26
B-3996kur $\Delta$ tdh $\Delta$ pckA/pVIC40	4.9	8.92

Example 8

Preparation of L-lysine with the strain TOC21R $\Delta$ pckA

- 5 The L-lysine-producing *E. coli* strain pDA1/TOC21R is described in the patent application F-A-2511032 and deposited at the Collection Nationale de Culture de Microorganisme (CNCM = National Microorganism Culture Collection, Pasteur Institute, Paris, France) under number  
10 I-167. The strain and the plasmid-free host are also described by Dauce-Le Reverend et al. (European Journal of Applied Microbiology and Biotechnology 15:227-231 (1982)) under the name TOC21R/pDA1.

- 8.1 Position-specific mutagenesis of the pckA gene in the  
15 *E. coli* strain TOC21R

After culture in antibiotic-free LB medium for approximately six generations, a derivative of strain pDA1/TOC21R which no longer contains the plasmid pDA1 is isolated. The strain formed is tetracycline-sensitive and  
20 is called TOC21R.

For replacement of the chromosomal pckA gene by the plasmid-coded deletion construct, TOC21R is transformed with the plasmid pMAK705 $\Delta$ pckA (Example 2). The gene replacement is carried out by the selection method  
25 described by Hamilton et al. (1989) Journal of Bacteriology

174, 4617 - 4622) and is verified by standard PCR methods (Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press) with the following oligonucleotide primers:

- 5 pckA'5'-1: 5' - GATCCGAGCCTGACAGGTTA - 3'  
pckA'3'-2: 5' - CCGGAGAAGCGTAGGTGTTA - 3'

The strain obtained is called TOC21RΔpckA.

## 8.2 Preparation of L-lysine with the strain TOC21RΔpckA

The formation of L-lysine by the strains TOC21RA pckA and  
10 TOC21R is checked in batch cultures of 10 ml contained in  
100 ml conical flasks. For this, 10 ml of preculture medium  
of the following composition: 2 g/l yeast extract, 10 g/l  
(NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 1 g/l KH<sub>2</sub>PO<sub>4</sub>, 0.5 g/l MgSO<sub>4</sub>\*7H<sub>2</sub>O, 15 g/l CaCO<sub>3</sub>, 20  
g/l glucose are inoculated and the batch is incubated for  
15 16 hours at 37°C and 180 rpm on an ESR incubator from  
Kühner AG (Birsfelden, Switzerland). 250 µl of this  
preculture are transinoculated into 10 ml of production  
medium (25 g/l (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 2 g/l KH<sub>2</sub>PO<sub>4</sub>, 1 g/l MgSO<sub>4</sub>\*7H<sub>2</sub>O,  
0.03 g/l FeSO<sub>4</sub>\*7H<sub>2</sub>O, 0.018 g/l MnSO<sub>4</sub>\*1H<sub>2</sub>O, 30 g/l CaCO<sub>3</sub>, 20  
20 g/l glucose, 25 mg/l L-isoleucine and 5 mg/l thiamine) and  
the batch is incubated for 72 hours at 37°C. After the  
incubation the optical density (OD) of the culture  
suspension is determined with an LP2W photometer from Dr.  
Lange (Berlin, Germany) at a measurement wavelength of 660  
25 nm.

The concentration of L-lysine formed is then determined in  
the sterile-filtered culture supernatant with an amino acid  
analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion  
exchange chromatography and post-column reaction with  
30 ninhydrin detection.

The result of the experiment is shown in Table 5.

Table 5

Strain	OD (660 nm)	L-Lysine g/l
TOC21R	1.0	1.14
TOC21R $\Delta$ pckA	1.0	1.27

Example 9

Preparation of L-isoleucine with the strain  
5 B-3996kur $\Delta$ t<sub>dh</sub>ilvA<sup>+</sup> $\Delta$ pckA/pVIC40

9.1 Preparation of the strain B-  
3996kur $\Delta$ t<sub>dh</sub>ilvA<sup>+</sup> $\Delta$ pckA/pVIC40

The strain B-3996kur $\Delta$ t<sub>dh</sub>, which is in need of L-isoleucin,  
obtained in Example 7.1 is transduced with the aid of the  
10 phage Plkc (Lennox, Virology 1, 190-206 (1955); Miller,  
Experiments in Molecular Genetics, Cold Spring Harbor  
Laboratory 1972) and L-isoleucine-prototrophic  
transductants are isolated.

For this, the phage Plkc is multiplied on the strain MG1655  
15 (Guyer et al., Cold Spring Harbor Symposium of Quantitative  
Biology 45, 135-140 (1981) and Blattner et al., Science  
277, 1453-1462 (1997)) and the phage lysate is employed for  
the transduction of the strain B-3996kur $\Delta$ t<sub>dh</sub>. The  
multiplicity of the infection is approximately 0.2.  
20 Selection for L-isoleucine-prototrophic transductants is  
carried out on minimal agar, which contains 2 g/l glucose  
and 10 mg/l L-threonine. An L-isoleucine-prototrophic  
transductant is isolated, smeared on to LB agar for  
purification or isolation and called B-3996kur $\Delta$ t<sub>dh</sub>ilvA<sup>+</sup>.

The pckA gene of the strain B-3996kur $\Delta$ tdhilvA<sup>+</sup> is then replaced, as described in Example 3, by the  $\Delta$ pckA allele prepared in Example 1 and 2. The strain obtained is called B-3996kur $\Delta$ tdhilvA<sup>+</sup> $\Delta$ pckA.

- 5 The strains B-3996kur $\Delta$ tdhilvA<sup>+</sup> and B-3996kur $\Delta$ tdhilvA<sup>+</sup> $\Delta$ pckA are transformed with the plasmid pVIC40 isolated from strain B-3996 and plasmid-carrying cells are selected on LB agar, which is supplemented with 20  $\mu$ g/ml streptomycin. In each case a selected individual colony is called B-  
10 3996kur $\Delta$ tdhilvA<sup>+</sup> $\Delta$ pckA/pVIC40 and B-3996kur $\Delta$ tdhilvA<sup>+</sup>/pVIC40.

## 9.2 Preparation of L-isoleucine

- The preparation of L-isoleucine by the strains B-3996kur $\Delta$ tdhilvA<sup>+</sup>/pVIC40 and B-3996kur $\Delta$ tdhilvA<sup>+</sup> $\Delta$ pckA/pVIC40 is tested under the test  
15 conditions as described in Example 4. The minimal medium, the preculture medium and the production medium are additionally supplemented with 20  $\mu$ g/ml streptomycin.

The result of the experiment is shown in Table 6.

Table 6

Strain	OD (660 nm)	L-Isoleucine mg/l
B-3996kur $\Delta$ tdhilvA <sup>+</sup> /pVIC40	5.8	57
B-3996kur $\Delta$ tdhilvA <sup>+</sup> $\Delta$ pckA/pVIC40	5.7	70

Example 10

Preparation of L-valine with the strain B-12288 $\Delta$ pckA

The L-valine-producing E. coli strain AJ 11502 is described in the patent specification US-A-4391907 and deposited at  
5 the National Center for Agricultural Utilization Research (Peoria, Illinois, USA) as NRRL B-12288.

10.1 Position-specific mutagenesis of the pckA gene in the E. coli strain B-1288

After culture in antibiotic-free LB medium for  
10 approximately six generations, a plasmid-free derivative of strain AJ 11502 is isolated. The strain formed is ampicillin-sensitive and is called AJ11502kur.

For replacement of the chromosomal pckA gene by the plasmid-coded deletion construct, AJ11502kur is transformed  
15 with the plasmid pMAK705 $\Delta$ pckA (see Example 2). The gene replacement is carried out by the selection method described by Hamilton et al. (1989) Journal of Bacteriology 174, 4617 - 4622) and is verified by standard PCR methods (Innis et al. (1990) PCR Protocols. A Guide to Methods and  
20 Applications, Academic Press) with the following oligonucleotide primers:

pckA'5'-1: 5' - GATCCGAGCCTGACAGGTTA - 3'

pckA'3'-2: 5' - CCGGAGAAGCGTAGGTGTTA - 3'

The strain obtained is called AJ11502kur $\Delta$ pckA. The plasmid  
25 described in the patent specification US-A-4391907, which carries the genetic information in respect of valine production, is isolated from strain NRRL B-12288. The strain AJ11502kur $\Delta$ pckA is transformed with this plasmid. One of the transformants obtained is called B-12288 $\Delta$ pckA.



10.2 Preparation of L-valine with the strain B-12288 $\Delta$ pckA

The formation of L-valine by the strains B-12288 $\Delta$ pckA and NRRL B-12288 is checked in batch cultures of 10 ml contained in 100 ml conical flasks. For this, 10 ml of  
5 preculture medium of the following composition: 2 g/l yeast extract, 10 g/l  $(\text{NH}_4)_2\text{SO}_4$ , 1 g/l  $\text{KH}_2\text{PO}_4$ , 0.5 g/l  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 15 g/l  $\text{CaCO}_3$ , 20 g/l glucose and 50 mg/l ampicillin are inoculated and the batch is incubated for 16 hours at 37°C and 180 rpm on an ESR incubator from Kühner AG (Birsfelden,  
10 Switzerland). 250  $\mu\text{l}$  of this preculture are transinoculated into 10 ml of production medium (25 g/l  $(\text{NH}_4)_2\text{SO}_4$ , 2 g/l  $\text{KH}_2\text{PO}_4$ , 1 g/l  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.03 g/l  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.018 g/l  $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ , 30 g/l  $\text{CaCO}_3$ , 20 g/l glucose, 5 mg/l thiamine and 50 mg/l ampicillin) and the batch is incubated for 72  
15 hours at 37°C. After the incubation the optical density (OD) of the culture suspension is determined with an LP2W photometer from Dr. Lange (Berlin, Germany) at a measurement wavelength of 660 nm.

The concentration of L-valine formed is then determined in  
20 the sterile-filtered culture supernatant with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in Table 7.

25

Table 7

Strain	OD (660 nm)	L-Valine g/l
NRRL B-12288	5.6	0.93
B-12288 $\Delta$ pckA	5.5	1.12

Example 11

Construction of deletion mutations of the ytfP-yjfa gene region

The ytfP-yjfa gene region is amplified from Escherichia coli K12 using the polymerase chain reaction (PCR) and synthetic oligonucleotides. Starting from the nucleotide sequence of the ytfP-yjfa gene region in E. coli K12 MG1655 (SEQ ID No. 5), the following PCR primers are synthesized (MWG Biotech, Ebersberg, Germany):

10 ytfP-1: 5' - GGCGATGTCGCAACAAGCTG - 3'  
ytfP-2: 5' - CTGTTTCATGGCCGCTTGCTG - 3'

The chromosomal E. coli K12 MG1655 DNA employed for the PCR is isolated according to the manufacturers instructions with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany).  
15 A DNA fragment approx. 1300 bp in size can be amplified with the specific primers under standard PCR conditions (Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press) with Taq-DNA polymerase (Gibco-BRL, Eggenstein, Germany). The PCR product is  
20 ligated with the vector pCR2.1TOPO (TOPO TA Cloning Kit, Invitrogen, Groningen, The Netherlands) in accordance with the manufacturers instructions and transformed into the E. coli strain TOP10F'. Selection of plasmid-carrying cells takes place on LB agar, to which 50 µg/ml ampicillin are  
25 added. After isolation of the plasmid DNA, successful cloning of the PCR product is checked with the restriction enzymes EcoRI and NsiI.

To generate a 337 bp deletion in the ytfP-yjfa region, the vector pCR2.1TOPOytfP-yjfa is cleaved with the restriction  
30 enzymes NdeI and SspI and the DNA fragment 4.8 kbp in size is ligated, after treatment with Klenow enzyme.

To generate a 90 bp deletion, the vector pCR2.1TOPOytfP-yjfA is cleaved with the enzymes NdeI and SplI and the DNA fragment 5 kbp in size is ligated, after treatment with Klenow enzyme.

- 5 The E. coli strain DH5 $\alpha$  is transformed with the ligation  
batches and plasmid-carrying cells are selected on LB agar,  
to which 50  $\mu$ g/ml ampicillin is added. After isolation of  
the plasmid DNA those plasmids in which the mutagenic DNA  
sequence shown in SEQ ID No. 6 and SEQ ID No. 7 is cloned  
10 are detected by control cleavage with the enzyme EcoRI. The  
plasmids are called pCR2.1TOPO $\Delta$ yjfA and pCR2.1TOPO $\Delta$ 90bp.

Example 12

Construction of the replacement vectors pMAK705ΔyjfA and pMAK705Δ90bp

The ytfP-yjfA alleles described in Example 11 are isolated  
5 from the vectors pCR2.1TOPOΔyjfA and pCR2.1TOPOΔ90bp after  
restriction with the enzymes SacI and XbaI and separation  
in 0.8% agarose gel, and ligated with the plasmid pMAK705  
(Hamilton et al. (1989) Journal of Bacteriology 174, 4617 -  
4622), which is digested with the enzymes SacI and XbaI.  
10 The ligation batches are transformed in DH5α and plasmid-  
carrying cells are selected on LB agar, to which 20 μg/ml  
chloramphenicol are added. Successful cloning is  
demonstrated after isolation of the plasmid DNA and  
cleavage with the enzymes SacI and XbaI. The replacement  
15 vectors formed, pMAK705ΔyjfA (= pMAK705deltayjfA) and  
pMAK705Δ90bp (= pMAK705delta90bp), are shown in Figure 2  
and in Figure 5.

Example 13

Position-specific mutagenesis of the ytfP-yjfA gene region  
20 in the E. coli strain MG442

For replacement of the chromosomal ytfP-yjfA gene region  
with the plasmid-coded 90 bp deletion construct, MG442 is  
transformed with the plasmid pMAK705Δ90bp. The gene  
replacement is carried out by the selection method  
25 described by Hamilton et al. (1989) Journal of Bacteriology  
174, 4617 - 4622) and is verified by standard PCR methods  
(Innis et al. (1990) PCR Protocols. A Guide to Methods and  
Applications, Academic Press) with the following  
oligonucleotide primers:

30 ytfP-1: 5' - GGCGATGTCGCAACAAGCTG - 3'  
ytfP-2: 5' - CTGTTTCATGGCCGCTTGCTG - 3'

The strain obtained is called MG442 $\Delta$ 90yjfA.

#### Example 14

Preparation of L-threonine with the strain MG442 $\Delta$ 90yjfA

5 The preparation of L-threonine by the strain MG442 $\Delta$ 90yjfA is tested as described in Example 4. The result of the experiment is summarized in Table 8.

Table 8

Strain	OD (660 nm)	L-Threonine g/l
MG442	6.0	1.5
MG442 $\Delta$ 90yjfA	5.7	2.1

#### Example 15

10 Preparation of L-threonine with the strain MG442 $\Delta$ 90yjfA $\Delta$ pckA

15.1 Preparation of the strain MG442 $\Delta$ 90yjfA $\Delta$ pckA

The pckA gene of the strain MG442 $\Delta$ 90yjfA is replaced, as described in Example 3, by the  $\Delta$ pckA allele (see Example 1  
15 and 2). The strain obtained is called MG442 $\Delta$ 90yjfA $\Delta$ pckA.

15.2 Preparation of L-threonine

The preparation of L-threonine with the strain MG442 $\Delta$ 90yjfA $\Delta$ pckA is carried out as described in Example 4. The result is shown in Table 9.

Table 9

Strain	OD (660 nm)	L-Threonine g/l
MG442Δ90yjfA	5.7	2.1
MG442Δ90yjfAΔpckA	5.3	3.9

## Brief Description of the Figures:

- Figure 1: pMAK705ΔpckA ( = pMAK705deltapckA)
- 5 • Figure 2: pMAK705ΔyjfA ( = pMAK705deltayjfA)
- Figure 3: pMW218gdhA
- Figure 4: pMW219rhtC
- Figure 5: pMAK705Δ90bp ( = pMAK705delta90bp)

10 The length data are to be understood as approx. data. The abbreviations and designations used have the following meaning:

- cat: Chloramphenicol resistance gene
- rep-ts: Temperature-sensitive replication region of the plasmid pSC101
- 15 • pck1: Part of the 5' region of the pckA gene
- pck2: Part of the 3' region of the pckA gene
- ytfP'-yjfA': DNA sequence containing truncated coding regions of ytfP and yjfA
- kan: Kanamycin resistance gene
- 20 • gdhA: Glutamate dehydrogenase gene

- rhtC: Threonine resistance-imparting gene

The abbreviations for the restriction enzymes have the following meaning

- 5 • BamHI: restriction endonuclease from *Bacillus amyloliquefaciens*
- BglII: restriction endonuclease from *Bacillus globigii*
- ClaI: restriction endonuclease from *Caryophanon latum*
- EcoRI: restriction endonuclease from *Escherichia coli*
- 10 • EcoRV: restriction endonuclease from *Escherichia coli*
- HindIII: restriction endonuclease from *Haemophilus influenzae*
- KpnI: restriction endonuclease from *Klebsiella pneumoniae*
- 15 • PstI: restriction endonuclease from *Providencia stuartii*
- PvuI: restriction endonuclease from *Proteus vulgaris*
- SacI: restriction endonuclease from *Streptomyces achromogenes*
- 20 • SalI: restriction endonuclease from *Streptomyces albus*
- SmaI: restriction endonuclease from *Serratia marcescens*
- 25 • XbaI: restriction endonuclease from *Xanthomonas badrii*

- XhoI: restriction endonuclease from *Xanthomonas holcicola*



**What is claimed is:**

1. Fermentation process for the preparation of L-amino acids, especially L-threonine, wherein the following steps are carried out:
  - 5       a) fermentation of the microorganisms of the family Enterobacteriaceae producing the desired L-amino acid, in which microorganisms at least the pckA gene or nucleotide sequences coding therefor are attenuated and, in particular, switched off,
  - 10       b) enrichment of the L-amino acid in the medium or in the bacterial cells, and
  - 15       c) isolation of the L-amino acid, constituents of the fermentation broth and the biomass in its entirety or portions thereof optionally being isolated as a solid product together with the L-amino acid.
2. Process according to claim 1, wherein microorganisms are used in which other genes of the biosynthetic pathway of the desired L-amino acid are additionally amplified.
- 20 3. Process according to claim 1, wherein microorganisms are used in which the metabolic pathways which reduce the formation of the desired L-amino acid are at least partially switched off.
- 25 4. Process according to claim 1, wherein the expression of the polynucleotide(s) coding for the pckA gene is attenuated and, in particular, switched off.
5. Process according to claim 1, wherein the regulatory and/or catalytic properties of the polypeptide (enzyme protein) coded for by the polynucleotide pckA are  
30 reduced.

6. Process according to claim 1, wherein microorganisms of the family Enterobacteriaceae in which one or more genes selected from the group comprising:

- 5           6.1       the thrABC operon coding for aspartate kinase, homoserine dehydrogenase, homoserine kinase and threonine synthase,
- 6.2       the pyc gene coding for pyruvate carboxylase,
- 6.3       the pps gene coding for phosphoenolpyruvate synthase,
- 10          6.4       the ppc gene coding for phosphoenolpyruvate carboxylase,
- 6.5       the pntA and pntB genes coding for transhydrogenase,
- 6.6       the rhtB gene for homoserine resistance, and
- 15          6.7       the rhtC gene for threonine resistance,
- 6.8       the gdhA gene coding for glutamate dehydrogenase

are simultaneously amplified and, in particular, overexpressed are fermented for the preparation of L-amino acids.

20

7. Process according to claim 1, wherein microorganisms of the family Enterobacteriaceae in which one or more genes selected from the group comprising:

- 25           7.1       the tdh gene coding for threonine dehydrogenase,
- 7.2       the mdh gene coding for malate dehydrogenase,

7.3 the gene product of the open reading frame  
(orf) yjfA, and

7.4 the gene product of the open reading frame  
(orf) ytfP,

5 are attenuated and, in particular, switched off, or the  
expression is reduced, are fermented for the  
preparation of L-amino acids.

8. Fermentation process for the preparation of L-amino  
acids, especially L-threonine, wherein the following  
10 steps are carried out:

a) fermentation of the microorganisms of the family  
Enterobacteriaceae producing the desired L-amino  
acid, in which microorganisms at least the open  
reading frames yjfA and/or ytfP or nucleotide  
15 sequences coding therefor are attenuated and, in  
particular, switched off,

b) enrichment of the L-amino acid in the medium or in  
the bacterial cells, and

c) isolation of the L-amino acid, constituents of the  
20 fermentation broth and the biomass in its entirety  
or portions thereof optionally being isolated as a  
solid product together with the L-amino acid.

9. Process according to claim 1 or 8, wherein L-  
isoleucine, L-valine, L-lysine or L-threonine is  
25 prepared.

10. L-Amino acid-producing microorganisms of the family  
Enterobacteriaceae in which at least the pckA gene or  
nucleotide sequences coding therefor are attenuated  
and, in particular, switched off.

11. L-Amino acid-producing microorganisms of the family Enterobacteriaceae according to claim 10, which additionally have one or more features selected from the group comprising: a resistance to  $\alpha$ -amino- $\beta$ -hydroxyvaleric acid, an amplified homoserine dehydrogenase I-aspartate kinase I in the feed back resistant form, an optionally compensable partial need for L-isoleucine, an attenuated threonine dehydrogenase and the ability to utilize sucrose.
12. L-Amino acid-producing microorganisms of the family Enterobacteriaceae, in which at least the open reading frame yjfA and/or ytfP or nucleotide sequences coding therefor are attenuated and, in particular, switched off.
13. L-Amino acid-producing microorganisms of the family Enterobacteriaceae according to claim 12, which additionally have one or more features selected from the group comprising: a resistance to  $\alpha$ -amino- $\beta$ -hydroxyvaleric acid, an amplified homoserine dehydrogenase I-aspartate kinase I in the feed back resistant form, an optionally compensable partial need for L-isoleucine, an attenuated threonine dehydrogenase and the ability to utilize sucrose.
14. Plasmid pMAK705 $\Delta$ pckA, shown in Figure 1, containing parts of the 5' and 3' regions of the pckA gene, corresponding to SEQ ID No. 3.
15. Plasmid pMAK705 $\Delta$ yjfA, shown in Figure 2, containing the 5' and 3' flanks of the ytfP-yjfA region, including very short residues of the open reading frames yjfA- and ytfP, corresponding to SEQ ID No. 6.
16. Plasmid pMAK705 $\Delta$ 90bp, shown in Figure 5, containing the 5' and 3' flanks of the ytfP-yjfA region, including very

short residues of the open reading frames yjfA- and ytfP, corresponding to SEQ ID No. 7.

17. Isolated polynucleotide from microorganisms of the family Enterobacteriaceae containing a polynucleotide sequence coding for the 5' and 3' regions of the pckA gene, shown in SEQ ID No. 4, which is particularly suitable as a constituent of plasmids for the position-specific mutagenesis of the pckA gene.
18. Isolated polynucleotide from microorganisms of the family Enterobacteriaceae containing the 5' and 3' flanks of the ytfP-yjfA region, shown in SEQ ID No. 6, which is particularly suitable as a constituent of plasmids for the position-specific mutagenesis of the open reading frames ytfP and/or yjfA.
19. L-Threonine-producing strains of the family Enterobacteriaceae containing a deletion mutation in the pckA gene, corresponding to SEQ ID No. 4.
20. L-Threonine-producing strains of the family Enterobacteriaceae containing a deletion mutation in the open reading frame ytfP, corresponding to SEQ ID No. 6 or 7.
21. L-Threonine-producing strains of the family Enterobacteriaceae containing a deletion mutation in the open reading frame yjfA, corresponding to SEQ ID No. 6 or 7.
22. L-Threonine-producing strains of the family Enterobacteriaceae according to claim 19, additionally containing a deletion mutation in the open reading frame ytfP, corresponding to SEQ ID No. 6 or 7.
23. L-Threonine-producing strains of the family Enterobacteriaceae according to claim 19, additionally

containing a deletion mutation in the open reading frame yjfA, corresponding to SEQ ID No. 6 or 7.

24. L-Threonine-producing strains of the family Enterobacteriaceae according to claims 19, 20 or 21, wherein they have one or more features selected from the group comprising: a resistance to  $\alpha$ -amino- $\beta$ -hydroxyvaleric acid, an amplified homoserine dehydrogenase I-aspartate kinase I in the feed back resistant form, an optionally compensable partial need for L-isoleucine, an attenuated threonine dehydrogenase and the ability to utilize sucrose.
25. *Escherichia coli* K-12 strain MG442 $\Delta$ pckA deposited under number DSM 13761 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (German Collection of Microorganisms and Cell Cultures).
26. *Escherichia coli* K-12 strain MG442 $\Delta$ 90yjfA deposited under number DSM 14289 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (German Collection of Microorganisms and Cell Cultures).
27. *Escherichia coli* K-12 strain B3996kur $\Delta$ tdhpckA/PVIC40, deposited under number DSM 14150 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (German Collection of Microorganisms and Cell Cultures).

Figure 1:




Figure 2:






Figure 3:




Figure 4:





Figure 5:



## SEQUENCE PROTOCOL

&lt;110&gt; Degussa AG

5 <120> Process for the fermentative preparation of  
L-amino acids using strains of the  
Enterobacteriaceae family.

10 &lt;130&gt; 000425 BT

&lt;140&gt;

&lt;141&gt;

15 &lt;160&gt; 19

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 1622

20 &lt;212&gt; DNA

&lt;213&gt; Escherichia coli

&lt;220&gt;

&lt;221&gt; CDS

25 &lt;222&gt; (1)..(1620)

&lt;223&gt; pckA

&lt;400&gt; 1

30 atg cgc gtt aac aat ggt ttg acc ccg caa gaa ctc gag gct tat ggt 48  
Met Arg Val Asn Asn Gly Leu Thr Pro Gln Glu Leu Glu Ala Tyr Gly  
1 5 10 15

35 atc agt gac gta cat gat atc gtt tac aac cca agc tac gac ctg ctg 96  
Ile Ser Asp Val His Asp Ile Val Tyr Asn Pro Ser Tyr Asp Leu Leu  
20 25 30

40 tat cag gaa gag ctc gat ccg agc ctg aca ggt tat gag cgc ggg gtg 144  
Tyr Gln Glu Glu Leu Asp Pro Ser Leu Thr Gly Tyr Glu Arg Gly Val  
35 40 45

45 tta act aat ctg ggt gcc gtt gcc gtc gat acc ggg atc ttc acc ggt 192  
Leu Thr Asn Leu Gly Ala Val Ala Val Asp Thr Gly Ile Phe Thr Gly  
50 55 60

50 cgt tca cca aaa gat aag tat atc gtc cgt gac gat acc act cgc gat 240  
Arg Ser Pro Lys Asp Lys Tyr Ile Val Arg Asp Asp Thr Thr Arg Asp  
65 70 75 80

50 act ttc tgg tgg gca gac aaa ggc aaa ggt aag aac gac aac aaa cct 288  
Thr Phe Trp Trp Ala Asp Lys Gly Lys Gly Lys Asn Asp Asn Lys Pro  
85 90 95

55 ctc tct ccg gaa acc tgg cag cat ctg aaa ggc ctg gtg acc agg cag 336  
Leu Ser Pro Glu Thr Trp Gln His Leu Lys Gly Leu Val Thr Arg Gln  
100 105 110

60 ctt tcc ggc aaa cgt ctg ttc gtt gtc gac gct ttc tgt ggt gcg aac 384  
Leu Ser Gly Lys Arg Leu Phe Val Val Asp Ala Phe Cys Gly Ala Asn  
115 120 125

	ccg gat act cgt ctt tcc gtc cgt ttc atc acc gaa gtg gcc tgg cag	432
	Pro Asp Thr Arg Leu Ser Val Arg Phe Ile Thr Glu Val Ala Trp Gln	
	130 135 140	
5	gcg cat ttt gtc aaa aac atg ttt att cgc ccg agc gat gaa gaa ctg	480
	Ala His Phe Val Lys Asn Met Phe Ile Arg Pro Ser Asp Glu Glu Leu	
	145 150 155 160	
10	gca ggt ttc aaa cca gac ttt atc gtt atg aac ggc gcg aag tgc act	528
	Ala Gly Phe Lys Pro Asp Phe Ile Val Met Asn Gly Ala Lys Cys Thr	
	165 170 175	
15	aac ccg cag tgg aaa gaa cag ggt ctc aac tcc gaa aac ttc gtg gcg	576
	Asn Pro Gln Trp Lys Glu Gln Gly Leu Asn Ser Glu Asn Phe Val Ala	
	180 185 190	
20	ttt aac ctg acc gag cgc atg cag ctg att ggc ggc acc tgg tac ggc	624
	Phe Asn Leu Thr Glu Arg Met Gln Leu Ile Gly Gly Thr Trp Tyr Gly	
	195 200 205	
25	ggc gaa atg aag aaa ggg atg ttc tcg atg atg aac tac ctg ctg ccg	672
	Gly Glu Met Lys Lys Gly Met Phe Ser Met Met Asn Tyr Leu Leu Pro	
	210 215 220	
30	ctg aaa ggt atc gct tct atg cac tgc tcc gcc aac gtt ggt gag aaa	720
	Leu Lys Gly Ile Ala Ser Met His Cys Ser Ala Asn Val Gly Glu Lys	
	225 230 235 240	
35	ggc gat gtt gcg gtg ttc ttc ggc ctt tcc ggc acc ggt aaa acc acc	768
	Gly Asp Val Ala Val Phe Phe Gly Leu Ser Gly Thr Gly Lys Thr Thr	
	245 250 255	
40	ctt tcc acc gac ccg aaa cgt cgc ctg att ggc gat gac gaa cac ggc	816
	Leu Ser Thr Asp Pro Lys Arg Arg Leu Ile Gly Asp Asp Glu His Gly	
	260 265 270	
45	tgg gac gat gac ggc gtg ttt aac ttc gaa ggc ggc tgc tac gca aaa	864
	Trp Asp Asp Asp Gly Val Phe Asn Phe Glu Gly Gly Cys Tyr Ala Lys	
	275 280 285	
50	act atc aag ctg tcg aaa gaa gcg gaa cct gaa atc tac aac gct atc	912
	Thr Ile Lys Leu Ser Lys Glu Ala Glu Pro Glu Ile Tyr Asn Ala Ile	
	290 295 300	
55	cgt cgt gat gcg ttg ctg gaa aac gtc acc gtg cgt gaa gat ggc act	960
	Arg Arg Asp Ala Leu Leu Glu Asn Val Thr Val Arg Glu Asp Gly Thr	
	305 310 315 320	
60	atc gac ttt gat gat ggt tca aaa acc gag aac acc cgc gtt tct tat	1008
	Ile Asp Phe Asp Asp Gly Ser Lys Thr Glu Asn Thr Arg Val Ser Tyr	
	325 330 335	
65	ccg atc tat cac atc gat aac att gtt aag ccg gtt tcc aaa gcg ggc	1056
	Pro Ile Tyr His Ile Asp Asn Ile Val Lys Pro Val Ser Lys Ala Gly	
	340 345 350	
70	cac gcg act aag gtt atc ttc ctg act gct gat gct ttc ggc gtg ttg	1104
	His Ala Thr Lys Val Ile Phe Leu Thr Ala Asp Ala Phe Gly Val Leu	
	355 360 365	

	ccg ccg gtt tct cgc ctg act gcc gat caa acc cag tat cac ttc ctc	1152
	Pro Pro Val Ser Arg Leu Thr Ala Asp Gln Thr Gln Tyr His Phe Leu	
	370 375 380	
5	tct ggc ttc acc gcc aaa ctg gcc ggt act gag cgt ggc atc acc gaa	1200
	Ser Gly Phe Thr Ala Lys Leu Ala Gly Thr Glu Arg Gly Ile Thr Glu	
	385 390 395 400	
10	ccg acg cca acc ttc tcc gct tgc ttc ggc gcg gca ttc ctg tcg ctg	1248
	Pro Thr Pro Thr Phe Ser Ala Cys Phe Gly Ala Ala Phe Leu Ser Leu	
	405 410 415	
15	cac ccg act cag tac gca gaa gtg ctg gtg aaa cgt atg cag gcg gcg	1296
	His Pro Thr Gln Tyr Ala Glu Val Leu Val Lys Arg Met Gln Ala Ala	
	420 425 430	
20	ggc gcg cag gct tat ctg gtt aac act ggc tgg aac ggc act ggc aaa	1344
	Gly Ala Gln Ala Tyr Leu Val Asn Thr Gly Trp Asn Gly Thr Gly Lys	
	435 440 445	
	cgt atc tcg att aaa gat acc cgc gcc att atc gac gcc atc ctc aac	1392
	Arg Ile Ser Ile Lys Asp Thr Arg Ala Ile Ile Asp Ala Ile Leu Asn	
	450 455 460	
25	ggg tcg ctg gat aat gca gaa acc ttc act ctg ccg atg ttt aac ctg	1440
	Gly Ser Leu Asp Asn Ala Glu Thr Phe Thr Leu Pro Met Phe Asn Leu	
	465 470 475 480	
30	gcg atc cca acc gaa ctg ccg ggc gta gac acg aag att ctc gat ccg	1488
	Ala Ile Pro Thr Glu Leu Pro Gly Val Asp Thr Lys Ile Leu Asp Pro	
	485 490 495	
35	cgt aac acc tac gct tct ccg gaa cag tgg cag gaa aaa gcc gaa acc	1536
	Arg Asn Thr Tyr Ala Ser Pro Glu Gln Trp Gln Glu Lys Ala Glu Thr	
	500 505 510	
40	ctg gcg aaa ctg ttt atc gac aac ttc gat aaa tac acc gac acc cct	1584
	Leu Ala Lys Leu Phe Ile Asp Asn Phe Asp Lys Tyr Thr Asp Thr Pro	
	515 520 525	
	gcg ggt gcc gcg ctg gta gcg gct ggt ccg aaa ctg taa	1623
	Ala Gly Ala Ala Leu Val Ala Ala Gly Pro Lys Leu	
	530 535 540	
45	<210> 2	
	<211> 540	
	<212> PRT	
	<213> Escherichia coli	
50	<400> 2	
	Met Arg Val Asn Asn Gly Leu Thr Pro Gln Glu Leu Glu Ala Tyr Gly	
	1 5 10 15	
55	Ile Ser Asp Val His Asp Ile Val Tyr Asn Pro Ser Tyr Asp Leu Leu	
	20 25 30	
60	Tyr Gln Glu Glu Leu Asp Pro Ser Leu Thr Gly Tyr Glu Arg Gly Val	
	35 40 45	

Leu Thr Asn Leu Gly Ala Val Ala Val Asp Thr Gly Ile Phe Thr Gly  
 50 55 60  
 5 Arg Ser Pro Lys Asp Lys Tyr Ile Val Arg Asp Asp Thr Thr Arg Asp  
 65 70 75 80  
 Thr Phe Trp Trp Ala Asp Lys Gly Lys Gly Lys Asn Asp Asn Lys Pro  
 85 90 95  
 10 Leu Ser Pro Glu Thr Trp Gln His Leu Lys Gly Leu Val Thr Arg Gln  
 100 105 110  
 Leu Ser Gly Lys Arg Leu Phe Val Val Asp Ala Phe Cys Gly Ala Asn  
 115 120 125  
 15 Pro Asp Thr Arg Leu Ser Val Arg Phe Ile Thr Glu Val Ala Trp Gln  
 130 135 140  
 20 Ala His Phe Val Lys Asn Met Phe Ile Arg Pro Ser Asp Glu Glu Leu  
 145 150 155 160  
 Ala Gly Phe Lys Pro Asp Phe Ile Val Met Asn Gly Ala Lys Cys Thr  
 165 170 175  
 25 Asn Pro Gln Trp Lys Glu Gln Gly Leu Asn Ser Glu Asn Phe Val Ala  
 180 185 190  
 Phe Asn Leu Thr Glu Arg Met Gln Leu Ile Gly Gly Thr Trp Tyr Gly  
 195 200 205  
 30 Gly Glu Met Lys Lys Gly Met Phe Ser Met Met Asn Tyr Leu Leu Pro  
 210 215 220  
 35 Leu Lys Gly Ile Ala Ser Met His Cys Ser Ala Asn Val Gly Glu Lys  
 225 230 235 240  
 Gly Asp Val Ala Val Phe Phe Gly Leu Ser Gly Thr Gly Lys Thr Thr  
 245 250 255  
 40 Leu Ser Thr Asp Pro Lys Arg Arg Leu Ile Gly Asp Asp Glu His Gly  
 260 265 270  
 Trp Asp Asp Asp Gly Val Phe Asn Phe Glu Gly Gly Cys Tyr Ala Lys  
 275 280 285  
 45 Thr Ile Lys Leu Ser Lys Glu Ala Glu Pro Glu Ile Tyr Asn Ala Ile  
 290 295 300  
 50 Arg Arg Asp Ala Leu Leu Glu Asn Val Thr Val Arg Glu Asp Gly Thr  
 305 310 315 320  
 Ile Asp Phe Asp Asp Gly Ser Lys Thr Glu Asn Thr Arg Val Ser Tyr  
 325 330 335  
 55 Pro Ile Tyr His Ile Asp Asn Ile Val Lys Pro Val Ser Lys Ala Gly  
 340 345 350  
 His Ala Thr Lys Val Ile Phe Leu Thr Ala Asp Ala Phe Gly Val Leu  
 355 360 365  
 60

Pro Pro Val Ser Arg Leu Thr Ala Asp Gln Thr Gln Tyr His Phe Leu  
 370 375 380  
 5 Ser Gly Phe Thr Ala Lys Leu Ala Gly Thr Glu Arg Gly Ile Thr Glu  
 385 390 395 400  
 Pro Thr Pro Thr Phe Ser Ala Cys Phe Gly Ala Ala Phe Leu Ser Leu  
 405 410 415  
 10 His Pro Thr Gln Tyr Ala Glu Val Leu Val Lys Arg Met Gln Ala Ala  
 420 425 430  
 Gly Ala Gln Ala Tyr Leu Val Asn Thr Gly Trp Asn Gly Thr Gly Lys  
 435 440 445  
 15 Arg Ile Ser Ile Lys Asp Thr Arg Ala Ile Ile Asp Ala Ile Leu Asn  
 450 455 460  
 20 Gly Ser Leu Asp Asn Ala Glu Thr Phe Thr Leu Pro Met Phe Asn Leu  
 465 470 475 480  
 Ala Ile Pro Thr Glu Leu Pro Gly Val Asp Thr Lys Ile Leu Asp Pro  
 485 490 495  
 25 Arg Asn Thr Tyr Ala Ser Pro Glu Gln Trp Gln Glu Lys Ala Glu Thr  
 500 505 510  
 Leu Ala Lys Leu Phe Ile Asp Asn Phe Asp Lys Tyr Thr Asp Thr Pro  
 515 520 525  
 30 Ala Gly Ala Ala Leu Val Ala Ala Gly Pro Lys Leu  
 530 535 540  
 35  
 <210> 3  
 <211> 1156  
 <212> DNA  
 <213> Escherichia coli  
 40  
 <220>  
 <221> misc\_feature  
 <222> (1)..(1156)  
 <223> Mutagene DNA  
 45  
 <220>  
 <221> misc\_feature  
 <222> (1)..(35)  
 <223> Technical DNA/residues of the polylinker sequence  
 50  
 <220>  
 <221> misc\_feature  
 <222> (36)..(522)  
 <223> Part of the 5' region (pck1) of the pckA gene  
 55  
 <220>  
 <221> misc\_feature  
 <222> (523)..(542)  
 <223> Technical DNA/residues of the polylinker sequence  
 60  
 <220>  
 <221> misc\_feature



```

<222> (543)..(1105)
<223> Part of the 3' region (pck2) of the pckA gene

<220>
5 <221> misc feature
  <222> (1106)..(1156)
  <223> Technical DNA/residues of the polylinker sequence

<400> 3
10 ctagtaacgg ccgccagtgt gctggaattc ggcttgatcc gagcctgaca ggttatgagc 60
   gcgggggtgtt aactaatctg ggtgccgttg ccgtcgatac cgggatcttc accggtcgtt 120
   caccaaaaga taagtatatc gtccgtgacg ataccactcg cgatactttc tgggtgggcag 180
   acaaaggcaa aggtaagaac gacaacaaac ctctctctcc ggaaacctgg cagcatctga 240
   aaggcctggt gaccaggcag ctttccggca aacgtctgtt cgttgctgac gctttctgtg 300
15 gtgcgaaccc ggatactcgt ctttccgtcc gtttcatcac cgaagtggcc tggcaggcgc 360
   attttgtcaa aaacatgttt attcgcccgga gcgatgaaga actggcaggt ttcaaaccag 420
   actttatcgt tatgaacggc gcgaagtgca ctaaccgcga gtggaaagaa cagggtctca 480
   actccgaaaa cttcgtggcg tttaacctga ccgagcgcac gcaagccgaa ttctgcagat 540
   cctgaagatg gcactatcga ctgtgatgat ggttcaaaaa ccgagaacac ccgcgtttct 600
20 tatccgatct atcacatcga taacattgtt aagccggttt ccaaagcggg ccacgcgact 660
   aaggttatct tcctgactgc tgatgctttc ggctgtgtgc cgcgcgtttc tcgcctgact 720
   gccgatcaaa ccagtatca cttcctctct ggcttcaccg ccaaactggc cgggtactgag 780
   cgtggcatca ccgaaccgac gccaaccttc tccgcttgct tcggcgcggc attcctgtcg 840
   ctgcaccgca ctcagtacgc agaagtgtgt gtgaaacgta tgcaggcggc gggcgcgcag 900
25 gcttatctgg ttaacactgg ctggaacggc actggcaaac gtatctcgat taaagatacc 960
   cgcgccatta tcgacgccat cctcaacggg tcgctggata atgcagaaac cttcactctg 1020
   ccgatgttta acctggcgat cccaaccgaa ctgccggggc tagacacgaa gattctcgat 1080
   ccgcgtaaca cctacgcttc tccggaagcc gaattctgca gatatccatc acactggcgg 1140
30 ccgctcgagc atgcat                                     1156

<210> 4
<211> 1294
<212> DNA
35 <213> Escherichia coli

<220>
<221> misc feature
<222> (1)..(3)
40 <223> Start codon of the delta pckA allele

<220>
<221> misc feature
<222> (1)..(598)
45 <223> 5' region of the delta pckA allele

<220>
<221> misc feature
<222> (599)..(618)
50 <223> Technical DNA/residues of the polylinker sequence

<220>
<221> misc feature
<222> (619)..(1291)
55 <223> 3' region of the delta pckA allele

<220>
<221> misc feature
<222> (1292)..(1294)
60 <223> Stop codon of the delta pckA allele

<400> 4

```

	atgcgcgtta	acaatggttt	gaccccgcaa	gaactcgagg	cttatggtat	cagtgcgta	60
	catgatatcg	tttacaaccc	aagctacgac	ctgctgtatc	aggaagagct	cgatccgagc	120
	ctgacagggt	atgagcgcg	ggtgttaact	aatctgggtg	ccgttgccgt	cgataccggg	180
	atcttcaccg	gtcgttcacc	aaaagataag	tatatcgccc	gtgacgatac	cactcgcgat	240
5	actttctggt	gggcagacaa	aggcaaagg	aagaacgaca	acaaacctct	ctctccggaa	300
	acctggcagc	atctgaaagg	cctggtgacc	aggcagcttt	ccggcaaacg	tctgttcggt	360
	gtcgacgctt	tctgtggtgc	gaacccggat	actcgtcttt	ccgtccggtt	catcaccgaa	420
	gtggcctggc	aggcgcatth	tgtcaaaaac	atgtttattc	gcccagagcg	tgaagaactg	480
	gcagggtttca	aaccagactt	tatcgttatg	aacggcgcg	agtgcactaa	cccgcagtgg	540
10	aaagaacagg	gtctcaactc	cgaaaacttc	gtggcggtta	acctgaccga	gcgcatgcaa	600
	gccgaattct	gcagatcctg	aagatggcac	tatcgacttt	gatgatggtt	caaaaaccga	660
	gaacacccgc	gtttcttatc	cgatctatca	catcgataac	attgttaagc	cggtttccaa	720
	agcggggccac	gcgactaagg	ttatcttcct	gactgctgat	gctttcggcg	tggtgcccgc	780
	ggtttctcgc	ctgactgccg	atcaaaccga	gtatcacttc	ctctctgggt	tcaccgccaa	840
15	actggccggg	actgagcgtg	gcataccgca	accgacgcca	accttctccg	cttgcttcgg	900
	cgcgccattc	ctgtcgctgc	acccgactca	gtacgcagaa	gtgctggtga	aacgtatgca	960
	ggcggcgggc	gcgcaggctt	atctggttaa	cactggctgg	aacggcactg	gcaaacgtat	1020
	ctcgattaaa	gatacccgcg	ccattatcga	cgccatcctc	aacggttcgc	tggtataatgc	1080
	agaaaccttc	actctgccga	tgtttaacct	ggcgatccca	accgaactgc	cgggcgtaga	1140
20	cacgaagatt	ctcgatccgc	gtaacaccta	cgcttctccg	gaacagtggc	aggaaaaagc	1200
	cgaaaccctg	gcgaaactgt	ttatcgacaa	cttcgataaa	tacaccgaca	cccctgcggg	1260
	tgcgcgctg	gtagcggtg	gtccgaaact	gtaa			1294
25	<210> 5						
	<211> 1248						
	<212> DNA						
	<213> Escherichia coli						
30	<220>						
	<221> gene						
	<222> (376)..(714)						
	<223> ORF ytfP						
35	<220>						
	<221> gene						
	<222> Complement((461)..(727))						
	<223> ORF yjfa						
40	<400> 5						
	ggcgatgtcg	caacaagctg	ccttgtctta	tttgetacgt	ggacaagggc	tggaagagcg	60
	tcagagcgac	agtgcggcaa	tgacctcgat	gctgattggt	ttgggggttg	cgcaaagtgg	120
	ccagattgtg	ggtaaaatcg	gcgagacgtt	tgccgtaagc	aatttagcgc	tcgacacca	180
	gggagtaggc	gactcctccc	aggtagtggt	cagcggctat	gtattgccag	gtctgcaagt	240
45	gaaatacggc	gtgggtatat	ttgactctat	agcaacactc	acgttacgtt	atcgccgtat	300
	gcctaagcta	tatctggaag	ccgtgtctgg	tgtagaccag	gcactggatt	tgctctatca	360
	gttcgagttt	tagcaatgcg	aatatttgtc	tacggcagtt	tacgccacaa	acaaggcaac	420
	agtcactgga	tgaccaatgc	ccagttactg	ggcgatttca	gtatcgataa	ctaccagttg	480
	tatagcctgg	gccactatcc	aggcgagttt	ccggggaaacg	gaacggtaca	cggtgaagtt	540
50	tatcgtattg	acaacgccac	gctggccgaa	cttgatgcct	tgccgaccag	gggcggtgaa	600
	tacgcgcgcc	agttgattca	gacgccgtac	gggagtgcat	ggatgtacgt	ttatcaacga	660
	cccgtcgatg	gattaaagct	aattgaaagc	ggcgactggt	tagacagggg	taagtaacca	720
	tatgcatacg	ccaccttcgg	gtggcggttg	tttttgcgag	acgactcgca	ttctgttttg	780
	taattccctc	accttttgc	ttctctccg	agccgctttc	catatctatt	aacgcataaa	840
55	aaactctgct	ggcattcaca	aatgcgcagg	ggtaaaacgt	ttcctgtagc	accgtgagtt	900
	atactttgta	taacttaagg	agggtcgagat	gcgtattacc	ataaaaagat	gggggaacag	960
	tgaggtatg	gtcattccca	atatcgtaat	gaaagaactt	aacttacagc	cggggcagag	1020
	cgtggagggc	caagtgaagc	acaatcaact	gattctgaca	cccattctca	ggcgctactc	1080
	gcttgatgaa	ctgctggcac	agtgtgacat	gaacgcgcgc	gaacttagcg	agcaggtagt	1140
60	ctggggtaaa	tccacccctg	cggggtgacga	aatatggtaa	agaaaagtga	atttgaaacg	1200
	ggagacattg	tgctgggtgg	ctttgatcca	gcaagcggcc	atgaacag		1248

```

    <210> 6
    <211> 911
    <212> DNA
5    <213> Escherichia coli

    <220>
    <221> misc_feature
    <222> (1)..(911)
10   <223> Deletion-carrying ytfP-yjfa region

    <220>
    <221> misc_feature
    <222> (1)..(383)
15   <223> 5' flank of the ytfP-yjfa region

    <220>
    <221> misc_feature
    <222> (384)..(911)
20   <223> 3' flank of the ytfP-yjfa region

    <220>
    <221> misc_feature
    <222> (376)..(378)
25   <223> ATG codon of the truncated ORF ytfP

    <220>
    <221> misc_feature
    <222> Complement((388)..(390))
30   <223> ATG codon of the truncated ORF yjfa

    <400> 6
    ggcgatgtcg caacaagctg ccttgtctta tttgctacgt ggacaagggc tggagagcga 60
    tcagagcgac agtgcggaac tgacctcgat gctgattggt ttggggggtg cgcaaagtgg 120
35   ccagattgtg ggtaaaatcg gcgagacgtt tggcgtaagc aatttagcgc tcgacaccca 180
    gggagtaggc gactcctccc aggtagtggg cagcggctat gtattgccag gtctgcaagt 240
    gaaatacggc gtgggtatat ttgactctat agcaacactc acgttacgtt atcgccctgat 300
    gcctaagcta tatctggaag ccgtgtctgg tgtagaccag gcaactggatt tgctctatca 360
    gttcgagttt tagcaatgcg aattatgcat acgccacctt cgggtggcgt tgttttttgc 420
40   gagacgactc gcattctggt ttgtaattcc ctcacctttt gcttttctct ccgagccgct 480
    ttccatatct attaacgcat aaaaaactct gctggcattc acaaatgcgc aggggtaaaa 540
    cgtttcctgt agcaccgtga gttatacttt gtataactta aggaggtgca gatgcgtatt 600
    accataaaaa gatgggggaa cagtgcaggt atggtcattc ccaatatcgt aatgaaagaa 660
    cttaacttac agccggggca gagcgtggag gcgcaagtga gcaacaatca actgattctg 720
45   acacccatct ccaggcgcta ctcgcttgat gaactgctgg cacagtgtga catgaacgcc 780
    gcggaactta gcgagcagga tgtctggggg aaatccaccc ctgcgggtga cgaaatatgg 840
    taaagaaaag tgaatttgaa cggggagaca ttgtgctggt tggctttgat ccagcaagcg 900
    gccatgaaca g
    911

50   <210> 7
    <211> 1158
    <212> DNA
    <213> Escherichia coli

55   <220>
    <221> misc_feature
    <222> (1)..(1158)
    <223> Deletion-carrying ytfP-yjfa region

60   <220>
    <221> misc_feature

```

```

<222> (1)..(630)
<223> 5' flank of the ytfP-yjfa region

<220>
5 <221> misc_feature
<222> (631)..(1158)
<223> 3' flank of the ytfP-yjfa region

<220>
10 <221> misc_feature
<222> (376)..(378)
<223> ATG codon of the truncated ORF ytfP

<220>
15 <221> misc_feature
<222> Complement((635)..(637))
<223> ATG codon of the truncated ORF yjfa

<400> 7
20 ggcgatgtcg caacaagctg ccttgtctta tttgctacgt ggacaagggc tggagagcga 60
   tcagagcgac agtgcggcaa tgacctcgat gctgattggt ttggggggtg cgcaaagtgg 120
   ccagattgtg ggtaaaatcg gcgagacgtt tggcgtaagc aatttagcgc tcgacaccca 180
   gggagtaggc gactcctccc aggtagtggg cagcggctat gtattgccag gtctgcaagt 240
   gaaatacggc gtgggtatat ttgactctat agcaaacctc acgttacgtt atcgctgat 300
25 gcctaagcta tatctggaag ccgtgtctgg tgtagaccag gcaactggatt tgctctatca 360
   gttcgagttt tagcaatgcg aatatttgtc tacggcagtt tacgccacaa acaaggcaac 420
   agtcactgga tgaccaatgc ccagttactg ggcgatttca gtatcgataa ctaccagttg 480
   tatagcctgg gccactatcc aggcgcagtt cgggggaacg gaacggtaca cgggtgaagt 540
   tatcgatttg acaacgccac gctggccgaa cttgatgcct tgcgcaccag gggcggtgaa 600
30 tacgcgcgcc agttgattca gacgccgtac tatgcatacg ccaccttcgg gtggcggttg 660
   tttttgcgag acgactcgca ttctgttttg taattccctc accttttgct tttctctccg 720
   agccgctttc catatctatt aacgcataaa aaactctgct ggcatcaca aatgcgcagg 780
   ggtaaaacgt ttcctgtagc accgtgagtt atactttgta taacttaagg aggtgcagat 840
   gcgtattacc ataaaaagat ggggggaacg tgcaggtatg gtcattccca atatcgtaat 900
35 gaaagaactt aacttacagc cggggcagag cgtggaggcg caagtgagca acaatcaact 960
   gattctgaca cccatctcca ggcgctactc gcttgatgaa ctgctggcac agtgtgacat 1020
   gaacgcgcgc gaacttagcg agcaggatgt ctggggtaaa tccaccctg cgggtgacga 1080
   aatatggtaa agaaaagtga atttgaacgg ggagacattg tgctggttgg ctttgatcca 1140
40 gcaagcggcc atgaacag                                     1158

<210> 8
<211> 20
<212> DNA
45 <213> Artificial sequence

<220>
<223> Description of the artificial sequence: Primer
      pckA'5'-1
50 <400> 8
   gatccgagcc tgacaggtta                                     20

<210> 9
<211> 20
<212> DNA
<213> Artificial sequence

60 <220>
<223> Description of the artificial sequence: Primer
      pckA'5'-2

```

	<400> 9		
	gcattgcgctc gggtcagggtta		20
5			
	<210> 10		
	<211> 22		
	<212> DNA		
10	<213> Artificial sequence		
	<220>		
	<223> Description of the artificial sequence: Primer		
	pckA'3'-1		
15	<400> 10		
	aggcctgaag atggcactat cg		22
20			
	<210> 11		
	<211> 20		
	<212> DNA		
	<213> Artificial sequence		
	<220>		
25	<223> Description of the artificial sequence: Primer		
	pckA'3'-2		
	<400> 11		
30	ccggagaagc gtaggtgtta		20
	<210> 12		
	<211> 20		
	<212> DNA		
35	<213> Artificial sequence		
	<220>		
	<223> Description of the artificial sequence: Primer Gdh1		
40	<400> 12		
	tgaacacttc tggcgggtacg		20
45			
	<210> 13		
	<211> 20		
	<212> DNA		
	<213> Artificial sequence		
	<220>		
50	<223> Description of the artificial sequence: Primer Gdh2		
	<400> 13		
	cctcggcgaa gctaatatgg		20
55			
	<210> 14		
	<211> 20		
	<212> DNA		
60	<213> Artificial sequence		
	<220>		
	<223> Description of the artificial sequence: Primer RhtC1		

	<400> 14	
	ctggttagcat cggcgaggca	20
5		
	<210> 15	
	<211> 20	
	<212> DNA	
	<213> Artificial sequence	
10		
	<220>	
	<223> Description of the artificial sequence: Primer RhtC2	
	<400> 15	
15	gcatgttgat ggcgatgacg	20
	<210> 16	
	<211> 20	
20	<212> DNA	
	<213> Artificial sequence	
	<220>	
	<223> Description of the artificial sequence: Primer Tdh1	
25		
	<400> 16	
	tcgcgaccta taagtgtggg	20
30	<210> 17	
	<211> 20	
	<212> DNA	
	<213> Artificial sequence	
35	<220>	
	<223> Description of the artificial sequence: Primer Tdh2	
	<400> 17	
40	aataccagcc cttgttcgtg	20
	<210> 18	
	<211> 20	
	<212> DNA	
45	<213> Artificial sequence	
	<220>	
	<223> Description of the artificial sequence: Primer	
	ytFP-1	
50		
	<400> 18	
	ggcgatgtcg caacaagctg	20
55	<210> 19	
	<211> 20	
	<212> DNA	
	<213> Artificial sequence	
60	<220>	
	<223> Description of the artificial sequence: Primer	
	ytFP-2	

<400> 19  
ctgttcacgg ccgcttgctg

20

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
11 April 2002 (11.04.2002)

PCT

(10) International Publication Number  
**WO 02/029080 A3**

(51) International Patent Classification<sup>7</sup>: **C12P 13/04**,  
13/08, C12N 1/21, 15/11, 15/60 // (C12P 13/04, C12R  
1:19) (C12P 13/08, C12R 1:19)

(21) International Application Number: PCT/EP01/10209

(22) International Filing Date:  
5 September 2001 (05.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
100 48 605.3 30 September 2000 (30.09.2000) DE  
100 55 516.0 9 November 2000 (09.11.2000) DE  
101 30 192.8 22 June 2001 (22.06.2001) DE

(71) Applicant: **DEGUSSA AG** [DE/DE]; Bennigsenplatz 1,  
40474 Düsseldorf (DE).

(72) Inventors: **RIEPING, Mechthild**; Mönkebergstrasse 1,  
33619 Bielefeld (DE). **BASTUCK, Christine**; Siekerwall  
17, 33602 Bielefeld (DE). **HERMANN, Thomas**; Zirkon-  
strasse 8, 33739 Bielefeld (DE). **THIERBACH, Georg**;  
Gunststrasse 21, 33613 Bielefeld (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,  
ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

**Published:**

— with international search report

(88) Date of publication of the international search report:  
26 September 2002

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



**WO 02/029080 A3**

(54) Title: FERMENTATION PROCESS FOR THE PREPARATION OF L-AMINO ACIDS USING STRAINS OF THE FAMILY ENTEROBACTERIACEAE

(57) Abstract: The invention relates to a fermentation process for the preparation of L-amino acids, especially L-threonine, in which the following steps are carried out: a) fermentation of the microorganisms of the family Enterobacteriaceae producing the desired L-amino acid, in which microorganisms at least the *pckA* gene and/or the open reading frames *yjfA* and *ytfP* are, individually or jointly, attenuated and, in particular, switched off, b) enrichment of the L-amino acid in the medium or in the bacterial cells, and c) isolation of the L-amino acid.



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/10209

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12P13/04 C12P13/08 C12N1/21 C12N15/11 C12N15/60  
 //(C12P13/04,C12R1:19),(C12P13/08,C12R1:19)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12P C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data, SEQUENCE SEARCH, EMBL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	EP 1 094 111 A (FORSCHUNGSZENTRUM JÜLICH GMBH) 25 April 2001 (2001-04-25)	1-7, 9-11,14, 17,19, 22-25,27
A	see especially claims 1-11 the whole document	8,9,12, 13,15, 16,18, 20,21, 24,26
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

30 April 2002

Date of mailing of the international search report

17.05.02

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Douschan, K

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/10209

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US; 1990  GOLDIE H ET AL: "PHYSICAL AND GENETIC  ANALYSIS OF THE PHOSPHOENOLPYRUVATE  CARBOXYKINASE PCK-A LOCUS FROM  ESCHERICHIA-COLI K12"  Database accession no. PREV199089092769  XP002193202  abstract  &amp; MOLECULAR &amp; GENERAL GENETICS,  vol. 220, no. 2, 1990, pages 191-196,  ISSN: 0026-8925</p> <p>---</p>	10
Y	<p>KRAMER R: "Genetic and physiological  approaches for the production of amino  acids"  JOURNAL OF BIOTECHNOLOGY, ELSEVIER SCIENCE  PUBLISHERS, AMSTERDAM, NL,  vol. 45, no. 1,  12 February 1996 (1996-02-12), pages 1-21,  XP004036833  ISSN: 0168-1656</p> <p>---</p>	1-7, 9-11,14, 17,19, 22-25,27
A	<p>see pages 3-5,13 and 15</p> <p>---</p>	8,9,12, 13,15, 16,18, 20,21, 24,26
Y	<p>DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US; 1990  MEDINA V ET AL: "SEQUENCE OF THE PCK-A  GENE OF ESCHERICHIA-COLI K-12 RELEVANCE TO  GENETIC AND ALLOSTERIC REGULATION AND  HOMOLOGY OF ESCHERICHIA-COLI  PHOSPHOENOLPYRUVATE CARBOXYKINASE WITH THE  ENZYMES FROM TRYPANOSOMA-BRUCI AND  SACCHAROMYCES-CEREVISIAE"  Database accession no. PREV199191038347  XP002193203  cited in the application  abstract  &amp; JOURNAL OF BACTERIOLOGY,  vol. 172, no. 12, 1990, pages 7151-7156,  ISSN: 0021-9193</p> <p>---</p> <p style="text-align: center;">-/--</p>	1-7, 9-11,14, 17,19, 22-25,27

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/10209

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US; March 1999 (1999-03)  PROST JEAN-FRANCOIS ET AL: "Detection of  an extended -10 element in the promoter  region of the pckA gene encoding  phosphoenolpyruvate carboxykinase in  Escherichia coli."  Database accession no. PREV199900300021  XP002193204  abstract  &amp; BIOCHIMIE (PARIS),  vol. 81, no. 3, March 1999 (1999-03),  pages 197-200,  ISSN: 0300-9084</p>	<p>1-7,  9-11,14,  17,19,  22-25,27</p>
A	<p>---  DATABASE CA 'Online!  CHEMICAL ABSTRACTS SERVICE, COLUMBUS,  OHIO, US;  RIVOLTA, CARLO ET AL: "A 35.7 kb DNA  fragment from the Bacillus subtilis  chromosome containing a putative 12.3 kb  operon involved in hexuronate catabolism  and a perfectly symmetrical hypothetical  catabolite-responsive element"  retrieved from STN  Database accession no. 129:63856  XP002197824  abstract  &amp; MICROBIOLOGY (READING, U. K.) (1998),  144(4), 877-884 ,</p>	<p>8,9,12,  13,15,  16,18,  20,21,  24,26</p>
A	<p>---  DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US; 1991  BLOMFIELD I C ET AL: "ALLELIC EXCHANGE IN  ESCHERICHIA-COLI USING THE  BACILLUS-SUBTILIS SAC-B GENE AND A  TEMPERATURE-SENSITIVE PSC-101 REPLICON"  Database accession no. PREV199192075325  XP002197825  abstract  &amp; MOLECULAR MICROBIOLOGY,  vol. 5, no. 6, 1991, pages 1447-1458,  ISSN: 0950-382X</p>	<p>15,16</p>

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP 01/10209

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-7, 9part., 10, 11, 14, 17, 19, 22, 23, 24part.,  
25, 27

Fermentation process for the preparation of L-amino acids using a microorganism of the family Enterobacteriaceae in which at least the pckA gene or nucleotide sequences coding therefore are attenuated; microorganisms useful therefore as well as plasmids and polynucleotides which suppress the pckA gene and PEP carboxykinase, respectively.

2. Claims: 8, 9part., 12, 13, 15, 16, 18, 20, 21, 24part., 26

Fermentation process for the preparation of L-amino acids using a microorganism of the family Enterobacteriaceae in which at least the open reading frames yjfA and/or ytfP or nucleotide sequences coding therefore are attenuated; microorganisms useful therefore as well as plasmids and polynucleotides which suppress the open reading frames yjfA and/or ytfP.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/10209

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1094111	A	25-04-2001	DE 19950409 A1	26-04-2001
			AU 6410400 A	26-04-2001
			BR 0004957 A	29-05-2001
			CN 1308125 A	15-08-2001
			EP 1094111 A2	25-04-2001
			JP 2001149086 A	05-06-2001
			PL 343339 A1	23-04-2001

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**